# antibodies -online.com







## anti-USP3 antibody (N-Term)

**Images** 



$\sim$						
	1//	Δ	r١	/1	<b>Θ</b> \	٨

Quantity:	0.4 mL
Target:	USP3
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This USP3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)
D	

#### **Product Details**

Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the N-terminal region of human USP3.
Isotype:	Ig Fraction
Specificity:	This antibody is specific to USP3 (N-term).
Purification:	Protein G Chromatography, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS.

## **Target Details**

Target:	USP3
Alternative Name:	USP3 (USP3 Products)

#### Target Details

Backo	round:
Backu	irouria.

Modification of target proteins by ubiquitin participates in a wide array of biological functions. Proteins destined for degradation or processing via the 26 S proteasome are coupled to multiple copies of ubiquitin. However, attachment of ubiquitin or ubiquitin-related molecules may also result in changes in subcellular distribution or modification of protein activity. An additional level of ubiquitin regulation, deubiquitination, is catalyzed by proteases called deubiquitinating enzymes, which fall into four distinct families. Ubiquitin C-terminal hydrolases, ubiquitin-specific processing proteases (USPs),1 OTU-domain ubiquitin-aldehyde-binding proteins, and Jab1/Pad1/MPN-domain-containing metallo-enzymes. Among these four families, USPs represent the most widespread and represented deubiquitinating enzymes across evolution. USPs tend to release ubiquitin from a conjugated protein. They display similar catalytic domains containing conserved Cys and His boxes but divergent N-terminal and occasionally C-terminal extensions, which are thought to function in substrate recognition, subcellular localization, and protein-protein interactions. Synonyms: Deubiquitinating enzyme 3, Ubiquitin carboxyl-terminal hydrolase 3, Ubiquitin thioesterase 3, Ubiquitin-specific-processing protease 3

Molecular Weight:	59097 Da
Gene ID:	9960, 5874
UniProt:	09Y6I4

### **Application Details**

Λ		NI - +
Ap	plication	Notes:

ELISA: 1/1,000. Western Blot: 1/100-1/500. Immunohistochemistry: 1/50-1/100.

Other applications not tested.

Optimal dilutions are dependent on conditions and should be determined by the user.

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS with 0.09 % (W/V) Sodium Azide as preservative.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Handling

Handling Advice:	Avoid repeated freezing and thawing.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at-20 °C for longer.	

## Images




Image 1.




Image 2.