

Datasheet for ABIN357572

anti-TSG101 antibody (N-Term)

2 Images



Go to Product page

| \sim | | | | |
|------------|---|----|-----------------|--|
| 0^{\vee} | 6 | rv | Ie _W | |

| Quantity: | 0.4 mL |
|----------------------|--|
| Target: | TSG101 |
| Binding Specificity: | N-Term |
| Reactivity: | Human, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This TSG101 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA) |
| Product Details | |
| Immunogen: | This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the N-terminal region of human TSG101. |
| Isotype: | lg Fraction |
| Specificity: | This antibody is specific to TSG101 (N-term). |
| Purification: | Protein G Chromatography, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS. |
| Target Details | |
| Target: | TSG101 |
| Alternative Name: | TSG101 (TSG101 Products) |
| | |

Target Details

| Background: | TSG101 belongs to a group of apparently inactive homologs of ubiquitin-conjugating enzymes. |
|---------------------|--|
| | The gene product contains a coiled-coil domain that interacts with stathmin, a cytosolic |
| | phosphoprotein implicated in tumorigenesis. The protein may play a role in cell growth and |
| | differentiation and act as a negative growth regulator. In vitro steady-state expression of this |
| | tumor susceptibility gene appears to be important for maintenance of genomic stability and cell |
| | cycle regulation. Mutations and alternative splicing in this gene occur in high frequency in |
| | breast cancer and suggest that defects occur during breast cancer tumorigenesis and/or |
| | progression. Synonyms: ESCRT-I complex subunit TSG101, Tumor susceptibility gene 101 |
| | protein |
| Molecular Weight: | 43944 Da |
| Gene ID: | 7251, 5874 |
| UniProt: | Q99816 |
| Application Details | |
| Application Notes: | ELISA: 1/1,000. Western Blot: 1/100-1/500. Immunohistochemistry: 1/50-1/100. |
| | Other applications not tested. |
| | Optimal dilutions are dependent on conditions and should be determined by the user. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 0.25 mg/mL |
| Buffer: | PBS with 0.09 % (W/V) Sodium Azide as preservative. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which |
| | should be handled by trained staff only. |
| Handling Advice: | Avoid repeated freezing and thawing. |
| Storage: | 4 °C/-20 °C |
| Storage Comment: | Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at-20 °C for longer. |
| | |




Image 1.




Image 2.