

## Datasheet for ABIN357599

## anti-UBE2E3 antibody (C-Term)





Go to Product page

$\sim$			
( )\	<b>/</b> e	rVI	iew

Quantity:	0.4 mL		
Target:	UBE2E3		
Binding Specificity:	C-Term		
Reactivity:	Human		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This UBE2E3 antibody is un-conjugated		
Application:	Western Blotting (WB), Enzyme Immunoassay (EIA)		
Product Details			
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide		
	selected from the C-terminal region of human UBCH9.		
Isotype:	Ig Fraction		
Specificity:	This antibody is specific to UBCH9/UBE2E3 (C-term). Predicted to cross react with Mouse (100		
	% Antigen Homology).		
Purification:	Protein G Chromatography, eluted with high and low pH buffers and neutralized immediately,		
	followed by dialysis against PBS.		
Target Details			
Target:			
rarget.	UBE2E3		

## Target Details

The modification of proteins with ubiquitin is an important cellular mechanism for targeting	
abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes	
of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and	
ubiquitin-protein ligases, or E3s. UBCH9 is a member of the E2 ubiquitin-conjugating enzyme	
family. The encoded protein shares 100 % sequence identity with the mouse and rat	
counterparts, which indicates that this enzyme is highly conserved in eukaryotes. Two	
alternatively spliced transcript variants encoding the same protein have been found. Synonyms:	
UBCE4, UBCH9, UbcM2, Ubiquitin carrier protein E3, Ubiquitin-conjugating enzyme E2 E3,	
Ubiquitin-conjugating enzyme E2-23 kDa, Ubiquitin-protein ligase E3	
22912 Da	
10477, 9606	
Q969T4	
ELISA: 1/1,000. Western Blot: 1/100-1/500.	
Other applications not tested.	
Optimal dilutions are dependent on conditions and should be determined by the user.	
For Research Use only	
Liquid	
0.25 mg/mL	
PBS with 0.09 % (W/V) Sodium Azide as preservative.	
Sodium azide	
This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
should be handled by trained staff only.	
Avoid repeated freezing and thawing.	
4 °C/-20 °C	
Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at-20 °C for longer.	




Image 1.