

Datasheet for ABIN357765

anti-ACOX1 antibody (C-Term)**1** Image[Go to Product page](#)

Overview

Quantity:	0.4 mL
Target:	ACOX1
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ACOX1 antibody is un-conjugated
Application:	Western Blotting (WB), Enzyme Immunoassay (EIA)

Product Details

Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the C-terminal region of human ACOX1.
Isotype:	Ig Fraction
Specificity:	This antibody is specific to ACOX1 (C-term).
Purification:	Protein G Chromatography, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS.

Target Details

Target:	ACOX1
Alternative Name:	ACOX1 (ACOX1 Products)

Target Details

Background:	ACOX1 is the first enzyme of the fatty acid beta-oxidation pathway, which catalyzes the desaturation of acyl-CoAs to 2-trans-enoyl-CoAs. It donates electrons directly to molecular oxygen, thereby producing hydrogen peroxide. Defects in this gene result in pseudoneonatal adrenoleukodystrophy, a disease that is characterized by accumulation of very long chain fatty acids. Synonyms: AOX, Palmitoyl-CoA oxidase, Peroxisomal acyl-coenzyme A oxidase 1, Straight-chain acyl-CoA oxidase
Molecular Weight:	74442 Da
Gene ID:	51, 9606
UniProt:	Q15067
Pathways:	Regulation of Lipid Metabolism by PPARalpha , Monocarboxylic Acid Catabolic Process

Application Details

Application Notes:	ELISA: 1/1,000. Western Blot: 1/50-1/100. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS with 0.09 % (W/V) Sodium Azide as preservative.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.




Image 1.