

Datasheet for ABIN357783 anti-PPT1 antibody (N-Term)

2 Images



Go to Product page

\sim		
()\//	erv	$\square \square \square$
\cup	_I V	$I \subset V \setminus$

Quantity:	0.4 mL
Target:	PPT1
Binding Specificity:	N-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PPT1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)
Product Details	
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the N-terminal region of human PPT1.
Isotype:	Ig Fraction
Specificity:	This antibody detects PPT1 (N-term).
Purification:	Protein G Chromatography eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS.
Target Details	
Target:	PPT1
Alternative Name:	PPT1 / PPT (PPT1 Products)

Target Details

Background:

Palmitoyl-protein thioesterase-1 (PPT1) is a lysosomal hydrolase that removes long-chain fatty acyl groups from modified cysteine residues in proteins. Mutations in PPT1 have been found to cause the infantile form of neuronal ceroid lipofuscinosis (INCL), and an animal model has been developed.1 The deduced PPT2 protein contains 302 amino acids, including a 27-amino acid leader peptide, a sequence motif characteristic of many thioesterases and lipases, and 5 potential N-linked glycosylation sites.2 PPT2 shares 18 % amino acid identity with PPT1.

Northern blot analysis detected a predominant 2.0-kb PPT2 transcript in the human tissues examined, with the highest expression in skeletal muscle, variable amounts of 2.8- and 7.0-kb transcripts were also observed. Recombinant PPT2, like PPT1, possesses thioesterase activity and localizes to the lysosome. Since PPT2 could not substitute for PPT1 in correcting the metabolic defect in INCL cells and was unable to remove palmitate groups from palmitoylated proteins that are routinely used as substrates for PPT1it has been postulated that PPT2 possesses a different substrate specificity than PPT1. Synonyms: PPT-1, Palmitoyl-protein hydrolase 1, Palmitoyl-protein thioesterase 1

Molecular Weight:	34193 Da
Gene ID:	5538, 5874
UniProt:	P50897
Pathways:	SARS-CoV-2 Protein Interactome

Application Details

Δnn	lication	Notes
, VDD	II Cation	110103

ELISA: 1/1,000. Western blot: 1/100-1/500. Immunohistochemistry: 1/50-1/100.

Other applications not tested.

Optimal dilutions are dependent on conditions and should be determined by the user.

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS with 0.09 % (W/V) Sodium Azide as preservative.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling

Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at-20 °C for longer.

Images




Image 1.




Image 2.