

Datasheet for ABIN357857  
**anti-PHPT1 antibody (N-Term)**



[Go to Product page](#)

2 Images

## Overview

Quantity:	0.4 mL
Target:	PHPT1
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PHPT1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)

## Product Details

Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the N-terminal region of human PHPT1.
Isotype:	Ig Fraction
Specificity:	This antibody detects PHPT1 (N-term). Predicted to cross react with Mouse and Rabbit (100 % Antigen Homology).
Purification:	Protein A Chromatography followed by peptide affinity purification.

## Target Details

Target:	PHPT1
Alternative Name:	PHPT1 ( <a href="#">PHPT1 Products</a> )

## Target Details

---

Background:	PHPT1 is an EDTA-insensitive phosphohistidine phosphatase that catalyzes the dephosphorylation of phosphopeptide I (Ek et al., 2002 [PubMed 12383260]). [supplied by OMIM]. Synonyms: 14 kDa phosphohistidine phosphatase, HSPC141, PHP14, Phosphohistidine phosphatase 1, Protein janus-A homolog
Molecular Weight:	13833 Da
Gene ID:	29085, 9606
UniProt:	<a href="#">Q9NRX4</a>
Pathways:	<a href="#">Positive Regulation of Peptide Hormone Secretion</a>

## Application Details

---

Application Notes:	ELISA: 1/1,000. Western Blot: 1/50-1/100. Immunohistochemistry: 1/10-1/50. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Restrictions:	For Research Use only

## Handling

---

Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS with 0.09 % (W/V) Sodium Azide as preservative.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.




Image 1.



Image 2.