antibodies - online.com







anti-TRAIP antibody (C-Term)

Images



Overview	
Quantity:	0.4 mL
Target:	TRAIP
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRAIP antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)
Product Details	
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the C-terminal region of human TRAIP.
Isotype:	Ig Fraction
Specificity:	This antibody detects TRAIP (C-term).
Purification:	Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Target Details	
Target:	TRAIP
Alternative Name:	TRAIP / RNF206 (TRAIP Products)

Target Details

rarget betails	
Background:	TRAIP is a protein that contains an N-terminal RING finger motif and a putative coiled-coil domain. A similar murine protein interacts with TNFR-associated factor 1 (TRAF1), TNFR-associated factor 2 (TRAF2), and cylindromatosis. The interaction with TRAF2 inhibits TRAF2 mediated nuclear factor kappa-B, subunit 1 activation that is required for cell activation and protection against apoptosis. Synonyms: RING finger protein 206, TRAF-interacting protein, TRIP
Molecular Weight:	53294 Da
Gene ID:	10293, 9606
UniProt:	Q9BWF2
Application Details	
Application Notes:	ELISA: 1/1,000. Western Blot: 1/50-1/100. Immunohistochemistry: 1/50-1/100. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS with 0.09 % (W/V) Sodium Azide as preservative.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at-20 °C for longer.




Image 1.




Image 2.