

## Datasheet for ABIN358703 anti-MMP 9 antibody (C-Term)

## 2 Images



Go to Product page

_						
	V	$\triangle$	r۱	/1	$\triangle$	Λ/
	' V '		ΙV			v v

Quantity:	0.4 mL	
Target:	MMP 9 (MMP9)	
Binding Specificity:	C-Term	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This MMP 9 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)	
Product Details		
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the C-terminal region of human MMP9.	
Isotype:	lg Fraction	
Specificity:	This antibody reacts to MMP9.	
Purification:	Protein G column, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS	
Target Details		
Target:	MMP 9 (MMP9)	
Alternative Name:	MMP-9 (MMP9 Products)	

## **Target Details**

_				
Background:	Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of			
	extracellular matrix in normal physiological processes, such as embryonic development,			
	reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and			
	metastasis. Most MMPs are secreted as inactive proproteins which are activated when cleaved			
	by extracellular proteinases. MMP9 degrades type IV and V collagens. Studies in rhesus			
	monkeys suggest that the enzyme is involved in IL-8-induced mobilization of hematopoietic			
	progenitor cells from bone marrow, and murine studies suggest a role in tumor-associated			
	tissue remodeling.Synonyms: 92 kDa gelatinase, 92 kDa type IV collagenase, CLG4B, GELB,			
	Gelatinase B, MMP9, Matrix metalloproteinase-9			
Gene ID:	4318, 9606			
UniProt:	P14780			
Pathways:	Cellular Response to Molecule of Bacterial Origin, Positive Regulation of Immune Effector			
	Process, CXCR4-mediated Signaling Events			
A 1: 1: D 1:				
Application Details				
Application Notes:	ELISA: 1/1,000. Western blotting: 1/100 - 1/500. Immunohistochemistry: 1/50 - 1/100.			
	Other applications not tested.			
	Optimal dilutions are dependent on conditions and should be determined by the user.			
Restrictions:	For Research Use only			
Handling				
Format:	Liquid			
Concentration:	0.25 mg/mL			
Buffer:	PBS with 0.09 % (W/V) sodium azide			
Preservative:	Sodium azide			
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which			
	should be handled by trained staff only.			
Handling Advice:	Avoid repeated freezing and thawing.			
Storage:	4 °C/-20 °C			
Storage Comment:	omment: Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at-20 °C for longer.			




Image 1.




Image 2.