antibodies .- online.com







anti-EPH Receptor A7 antibody (C-Term)

Images



\sim			
	N/6	1//r	$I \cap V$

Alternative Name:

Overview		
Quantity:	0.4 mL	
Target:	EPH Receptor A7 (EPHA7)	
Binding Specificity:	C-Term	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This EPH Receptor A7 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)	
Product Details		
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the C-terminal region of human EphA7.	
Isotype:	lg Fraction	
Specificity:	This antibody reacts to EphA7.	
Purification:	Protein G column, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS	
Target Details		
Target:	EPH Receptor A7 (EPHA7)	

EPHA7 (EPHA7 Products)


Target Details

Background:	Protein kinases are enzymes that transfer a phosphate group from a phosphate donor,	
	generally the g phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this	
	basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells,	
	regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement	
	and cell movement, apoptosis, and differentiation. With more than 500 gene products, the	
	protein kinase family is one of the largest families of proteins in eukaryotes. The family has	
	been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or	
	serine/threonine (STK) kinase catalytic domains. The tyrosine kinase (TK) group is mainly	
	involved in the regulation of cell-cell interactions such as differentiation, adhesion, motility and	
	death. There are currently about 90 TK genes sequenced, 58 are of receptor protein TK (e.g.	
	EGFR, EPH, FGFR, PDGFR, TRK, and VEGFR families), and 32 of cytosolic TK (e.g. ABL, FAK,	
	JAK, and SRC families).Synonyms: EHK3, EPH homology kinase 3, Ephrin type-A receptor 7,	
	HEK11, Receptor protein-tyrosine kinase HEK11, Tyrosine-protein kinase receptor EHK-3	
Gene ID:	2045, 9606	
UniProt:	Q15375	
Pathways:	RTK Signaling	
Application Details		
Application Notes:	ELISA: 1/1,000. Western blotting: 1/100 - 1/500. Immunohistochemistry: 1/50 - 1/100.	
	Other applications not tested.	
	Optimal dilutions are dependent on conditions and should be determined by the user.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	0.25 mg/mL	
Buffer:	PBS with 0.09 % (W/V) sodium azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Handling Advice:	Avoid repeated freezing and thawing.	
a.iaiiig / lavioc.	, word repeated treezing and thaming.	

Handling


Storage:	4 °C/-20 °C
Storage Comment: Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at-20 °C for l	

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining.



Western Blotting

Image 2. (LEFT)Western blot analysis of anti-EphA7 Pab in NCI-H460 cell lysate. EphA7 (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence. (RIGHT)Western blot analysis of EPHA7 (arrow) using EphA7 Antibody