

# Datasheet for ABIN359851

## anti-FGFR1 antibody (Tyr766)

## 1 Image



Go to Product page

	ve	rv	ie	W
$\circ$	v C	· I V	10	V V

Quantity:	0.4 mL
Target:	FGFR1
Binding Specificity:	Tyr766
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FGFR1 antibody is un-conjugated
Application:	Western Blotting (WB), Enzyme Immunoassay (EIA)
Product Details	
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide corresponding to amino acid residues surrounding Y766 of human FGFR1.
Immunogen:	
	corresponding to amino acid residues surrounding Y766 of human FGFR1.
Isotype:	corresponding to amino acid residues surrounding Y766 of human FGFR1.
Isotype: Specificity:	corresponding to amino acid residues surrounding Y766 of human FGFR1.  Ig Fraction  This antibody reacts to FGFR1.
Isotype: Specificity: Purification:	corresponding to amino acid residues surrounding Y766 of human FGFR1.  Ig Fraction  This antibody reacts to FGFR1.
Isotype: Specificity: Purification: Target Details	corresponding to amino acid residues surrounding Y766 of human FGFR1.  Ig Fraction  This antibody reacts to FGFR1.  Protein A column, followed by peptide affinity purification
Isotype: Specificity: Purification: Target Details Target:	corresponding to amino acid residues surrounding Y766 of human FGFR1.  Ig Fraction  This antibody reacts to FGFR1.  Protein A column, followed by peptide affinity purification  FGFR1

	is highly conserved between members and throughout evolution. FGFR family members differ	
	from one another in their ligand affinities and tissue distribution. A full-length representative	
	protein consists of an extracellular region, composed of three immunoglobulin-like domains, a	
	single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain.	
	The extracellular portion of the protein interacts with fibroblast growth factors, setting in	
	motion a cascade of downstream signals, ultimately influencing mitogenesis and	
	differentiation. This particular family member binds both acidic and basic fibroblast growth	
	factors and is involved in limb induction. Mutations in this gene can lead to Pfeiffer syndrome	
	and Jackson-Weiss syndrome. Synonyms: BFGFR, Basic fibroblast growth factor receptor 1,	
	CEK, FGFBR, FLG, FLT-2, FLT2, Fibroblast growth factor receptor 1, Fms-like tyrosine kinase 2,	
	HBGFR, N-sam, Proto-oncogene c-Fgr, bFGF-R-1	
Gene ID:	2260, 9606	
UniProt:	P11362	
Pathways:	RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin	

U Pa Signaling Pathway, Sensory Perception of Sound, Stem Cell Maintenance, S100 Proteins

### **Application Details**

Restrictions:	For Research Use only		
	Optimal dilutions are dependent on conditions and should be determined by the user.		
	Other applications not tested.		
Application Notes:	ELISA: 1/1,000. Western blotting: 1/50 - 1/100.		


### Handling

Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS with 0.09 % (W/V) sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C

Storage Comment:

Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at-20 °C for longer.

#### **Images**



#### **Western Blotting**

**Image 1.** Western blot analysis of FGFR1 (arrow) using rabbit polyclonal FGFR1 Antibody. 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the FGFR1 gene (Lane 2)