



[Go to Product page](#)

Datasheet for ABIN360552  
**anti-GRAF antibody (Middle Region)**

2 Images

Overview

Quantity:	0.4 mL
Target:	GRAF (ARHGAP26)
Binding Specificity:	Middle Region
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GRAF antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)

Product Details

Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the center region of human OPHN1L.
Isotype:	Ig Fraction
Specificity:	This antibody is specific to GRAF (OPHN1L).
Purification:	Protein G column, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS

Target Details

Target:	GRAF (ARHGAP26)
Alternative Name:	ARHGAP26 / OPHN1L ( <a href="#">ARHGAP26 Products</a> )

## Target Details

---

**Background:** Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the  $\gamma$  phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The STE group (homologs of yeast Sterile 7, 11, 20 kinases) consists of 50 kinases related to the mitogen-activated protein kinase (MAPK) cascade families (Ste7/MAP2K, Ste11/MAP3K, and Ste20/MAP4K). MAP kinase cascades, consisting of a MAPK and one or more upstream regulatory kinases (MAPKKs) have been best characterized in the yeast pheromone response pathway. Pheromones bind to Ste cell surface receptors and activate yeast MAPK pathway. Synonyms: GRAF, GTPase regulator associated with focal adhesion kinase, KIAA0621, Oligophrenin-1-like protein, Rho GTPase-activating protein 26, Rho-type GTPase-activating protein 26

---

**Gene ID:** 23092, 9606

---

**UniProt:** [Q9UNA1](#)

## Application Details

---

**Application Notes:** ELISA: 1/1,000. Western blotting: 1/100-1/500. Immunohistochemistry: 1/50-1/100.  
Other applications not tested.  
Optimal dilutions are dependent on conditions and should be determined by the user.

---

**Restrictions:** For Research Use only

## Handling

---

**Format:** Liquid

---

**Concentration:** 0.25 mg/mL

---

**Buffer:** PBS with 0.09 % (W/V) Sodium Azide as preservative.

---

**Preservative:** Sodium azide

---

**Precaution of Use:** This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

---

**Handling Advice:** Avoid repeated freezing and thawing.

---

## Handling

Storage: 4 °C/-20 °C

Storage Comment: Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at-20 °C for longer.

## Validation report #104331 for Multiplex Immunohistochemistry (mIHC)




Image 1.




Image 2.