# antibodies - online.com







# anti-PACSIN3 antibody (N-Term)

**Images** 



$\sim$						
	1//	Д	r۱	1	<b>Θ</b> 1	٨

Quantity:	0.4 mL	
Target:	PACSIN3	
Binding Specificity:	N-Term	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PACSIN3 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)	

#### **Product Details**

lmmunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the N-terminal region of human PACSIN3.
Isotype:	lg Fraction
Specificity:	This antibody reacts to PACSIN3.
Purification:	Protein G column, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS

# **Target Details**

Target:	PACSIN3	
Alternative Name:	PACSIN3 (PACSIN3 Products)	

## **Target Details**

Handling Advice:

Storage:

Larget Details	
Background:	Protein kinases are enzymes that transfer a phosphate group from a phosphate donor,
	generally the g phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this
	basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells,
	regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement
	and cell movement, apoptosis, and differentiation. With more than 500 gene products, the
	protein kinase family is one of the largest families of proteins in eukaryotes. The family has
	been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or
	serine/threonine (STK) kinase catalytic domains. The STE group (homologs of yeast Sterile 7,
	11, 20 kinases) consists of 50 kinases related to the mitogen-activated protein kinase (MAPK)
	cascade families (Ste7/MAP2K, Ste11/MAP3K, and Ste20/MAP4K). MAP kinase cascades,
	consisting of a MAPK and one or more upstream regulatory kinases (MAPKKs) have been best
	characterized in the yeast pheromone response pathway. Pheromones bind to Ste cell surface
	receptors and activate yeast MAPK pathway. Synonyms: Endophilin I, Protein kinase C and
	casein kinase substrate in neurons protein 3, SH3 domain-containing protein 6511
Gene ID:	29763, 9606
UniProt:	Q9UKS6
Application Details	
Application Notes:	ELISA: 1/1,000. Western blotting: 1/100 - 1/500. Immunohistochemistry: 1/50 - 1/100.
	Other applications not tested.
	Optimal dilutions are dependent on conditions and should be determined by the user.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS with 0.09 % (W/V) sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	along the department of the second se

should be handled by trained staff only.

Avoid repeated freezing and thawing.

4 °C/-20 °C

Storage Comment:

Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at-20 °C for longer.

## **Images**




Image 1.




Image 2.