antibodies - online.com







anti-E2F7 antibody (Internal Region)



Image



Overview

Quantity:	100 μg
Target:	E2F7
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This E2F7 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Purpose:	E2F7
Immunogen:	Peptide with sequence C-DLSKQKKFTPERNP, from the internal region of the protein sequence according to NP_976328.2.
Sequence:	DLSKQKKFTP ERNP
Isotype:	IgG
Cross-Reactivity:	Dog, Human, Pig
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

Target Details

Target Details	
Target:	E2F7
Alternative Name:	E2F7 (E2F7 Products)
Background:	E2F7, E2F transcription factor 7, FLJ12981
Gene ID:	144455
NCBI Accession:	NP_976328
Application Details	
Application Notes:	Western Blot: Approx 90 kDa band observed in Human Brain (Cerebellum) lysates (calculated MW of 99.9 kDa according to NP_976328.2). Recommended concentration: 0.3-1 µg/mL. Peptide ELISA: antibody detection limit dilution 1:32000.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

Western Blotting

Image 1. ABIN361188 (0.3μg/ml) staining of Human Cerebellum lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.