

### Datasheet for ABIN361188

# anti-E2F7 antibody (Internal Region)

## 1 Image



Go to Product page

0				

Quantity:	100 μg	
Target:	E2F7	
Binding Specificity:	Internal Region	
Reactivity:	Human	
Host:	Goat	
Clonality:	Polyclonal	
Conjugate:	This E2F7 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA	
Product Details		
Purpose:	E2F7	
	E2F7  Peptide with sequence C-DLSKQKKFTPERNP, from the internal region of the protein sequence according to NP_976328.2.	
Purpose:	Peptide with sequence C-DLSKQKKFTPERNP, from the internal region of the protein sequence	
Purpose: Immunogen:	Peptide with sequence C-DLSKQKKFTPERNP, from the internal region of the protein sequence according to NP_976328.2.	
Purpose: Immunogen: Sequence:	Peptide with sequence C-DLSKQKKFTPERNP, from the internal region of the protein sequence according to NP_976328.2.  DLSKQKKFTP ERNP	
Purpose: Immunogen: Sequence: Isotype:	Peptide with sequence C-DLSKQKKFTPERNP, from the internal region of the protein sequence according to NP_976328.2.  DLSKQKKFTP ERNP  IgG	

### **Target Details**

Target Details			
Target:	E2F7		
Alternative Name:	E2F7 (E2F7 Products)		
Background:	E2F7, E2F transcription factor 7, FLJ12981		
Gene ID:	144455		
NCBI Accession:	NP_976328		
Application Details			
Application Notes:	Western Blot: Approx 90 kDa band observed in Human Brain (Cerebellum) lysates (calculated MW of 99.9 kDa according to NP_976328.2). Recommended concentration: 0.3-1 µg/mL. Peptide ELISA: antibody detection limit dilution 1:32000.		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	0.5 mg/mL		
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.		
Handling Advice:	Minimize freezing and thawing.		
Storage:	-20 °C		
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.		

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

#### **Western Blotting**

**Image 1.** ABIN361188 (0.3μg/ml) staining of Human Cerebellum lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.