antibodies -online.com





anti-Peripherin antibody

3 Images



Publication



Go to Product page

Overview

Quantity:	100 μL
Target:	Peripherin (PRPH)
Reactivity:	Rat
Host:	Chicken
Clonality:	Polyclonal
Conjugate:	This Peripherin antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant rat peripherin expressed in and purified from E. coli
Specificity:	Specific for the ~57kDa peripherin protein.
Cross-Reactivity:	Human, Mouse (Murine), Rat (Rattus)
Predicted Reactivity:	most mammalian
Purification:	Total IgY fraction

Target Details

Target:	Peripherin (PRPH)
Alternative Name:	PRPH (PRPH Products)
Background:	Peripherin is a ~57kDa intermediate filament subunit found initially in sensory neurons of the
	peripheral nervous systems, which gives the protein its name. Subsequently, peripherin was

found in some sensory and other neurons of the central nervous system and also in PC12 cells. Peripherin is also expressed in certain neuroendocrine tumors and in the insulin producing cells of the pancreas. Peripherin belongs to the Class III family of intermediate filament subunits which also includes vimentin, glial fibrillary acidic protein (GFAP) and desmin. In contrast to the neurofilaments, peripherin is strongly up-regulated after nerve injury (1). Antibodies to peripherin can be used in identifying, classifying, and studying neurons throughout the nervous system. Peripherin is also a good diagnostic marker for ballooned axons seen in Lou Gehrig's disease (Amyotrophic lateral sclerosis) and some neuronally derived tumors (2). Autoantibodies to peripherin are frequently seen in the sera of patients with diabetes (3). Peripherin is not related to peripherin/RDS, a protein of the photoreceptor outer membrane mutations of which are causative of certain forms of slow retinal degeneration. Anti-Peripherin Western blot of rat cerebellar lysate showing specific immunolabeling of the ~ 57k peripherin protein.

Molecular Weight: '57 kDa

Gene ID: 3414

Application Details

Application Notes: Recommended Dilution: WB:1:1,000 IF: 1:500 Quality Control: Western blots performed on each lot.

Restrictions: For Research Use only

P21807

Handling

UniProt:

Format:	Liquid
Buffer:	total IgY fraction in PBS + 10 mM Sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C

Publications


Product cited in:

Sun, Wei, Xiong, Wang, Xie, Wang, Yang, Wang, Lu, Liu, Wang: "Synaptic released zinc promotes

tau hyperphosphorylation by inhibition of protein phosphatase 2A (PP2A)." in: **The Journal of biological chemistry**, Vol. 287, Issue 14, pp. 11174-82, (2012) (PubMed).


Chen, Xiong, Kong, Qu, Wang, Chen, Wang, Zhu: "Neuroglobin attenuates Alzheimer-like tau hyperphosphorylation by activating Akt signaling." in: **Journal of neurochemistry**, Vol. 120, Issue 1, pp. 157-64, (2011) (PubMed).

Validation report #103692 for Proximity Ligation Assay (PLA)




Western Blotting

Image 1. Western blots of rat cerebellar lysate showing specific immunolabeling of the \sim 57k peripherin protein.



Western Blotting

Image 2. Western blot of rat cerebellar lysate showing specific immunolabeling of the \sim 57 kDa peripherin protein.



Immunostaining

Image 3. Immunostaining of cultured rat neurons and glia showing peripherin in green and alpha-internexin in red. Cells expressing both peripherin and alpha-internexin appear yellow.