# antibodies -online.com





# anti-CAMKII gamma antibody

5 Images



Publication



Go to Product page

#### Overview

Quantity:	100 μg
Target:	CAMKII gamma (CAMK2G)
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CAMKII gamma antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunoprecipitation (IP), Immunofluorescence (IF), Immunocytochemistry (ICC), Radioimmunoassay (RIA)

#### **Product Details**

Immunogen:	Partially purified rat CaMKII
Clone:	6G9
Isotype:	lgG1
Specificity:	Detects ~50-60 kDa. Recognizes both phosphorylated and non-phosphorylated forms.
Cross-Reactivity:	Cow, Human, Mouse, Rat
Purification:	Protein G Purified

## **Target Details**

Target:	CAMKII gamma (CAMK2G)
Alternative Name:	CaMKII (CAMK2G Products)

## Target Details

Background:	CaMKII is an important member of the calcium/calmodulin-activated protein kinase family,
	functioning in neural synaptic stimulation and T-cell receptor signaling (1, 2). CaMKII is
	expressed in many different tissues but is specifically found in the neurons of the forebrain and
	its mRNA is found within the dendrites and the soma of the neuron. The CaMKII that is found in
	the neurons consist of two subunits of 52 (termed alpha genes) and 60 kDa (beta genes).
	CaMKII has catalytic and regulatory domains, as well as an ATP-binding domain, and a
	consensus phosphorylation site (3-7). The binding of Ca2+/calmodulin to its regulatory domain
	releases its auto inhibitory effect and activates the kinase (8). This kinase activation results in
	autophosphorylation at threonine 286 (8). The threonine phosphorylation state of CaMKII can
	be regulated through PP1/PKA. Whereas PP1 (protein phosphatase 1) dephosphorylates
	phospho-CaMKII at Thr286, PKA (protein kinase A) prevents this dephosphorylation (9).
	Autophosphorylation also enables CaMKII to attain an enhanced affinity for NMDA receptors in
	postsynaptic densities (10-12).
Gene ID:	12322
NCBI Accession:	NP_033922
UniProt:	P11798
Pathways:	WNT Signaling, Interferon-gamma Pathway, Hormone Transport, Myometrial Relaxation and
	Contraction, Regulation of long-term Neuronal Synaptic Plasticity
Application Details	
Application Notes:	• WB (1:10000)
	• IHC (1:2000)
	ICC/IF (1:50)      antimal dilutions for assays should be determined by the user.
	<ul> <li>optimal dilutions for assays should be determined by the user.</li> </ul>
Comment:	0.1 μg/ml was sufficient for detection of CamKII in 20 μg rat brain tissue extract by colorimetric
	immunoblot analysis using Goat Anti-Mouse IgG:AP as the secondary.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated

#### Handling


Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	-20°C

#### **Publications**

Product cited in:

Su, Gu, Wang, Wang: "Lidocaine attenuates proinflammatory cytokine production induced by extracellular adenosine triphosphate in cultured rat microglia." in: **Anesthesia and analgesia**, Vol. 111, Issue 3, pp. 768-74, (2010) (PubMed).

#### **Images**




#### **Immunohistochemistry**

Image 1. Immunohistochemistry analysis using Mouse Anti-CaMKII Monoclonal Antibody, Clone 6G9 (ABIN361643 and ABIN361644). Tissue: colon carcinoma. Species: Human. Fixation: Formalin. Primary Antibody: Mouse Anti-CaMKII Monoclonal Antibody (ABIN361643 and ABIN361644) at 1:10000 for 12 hours at 4 °C. Secondary Antibody: Biotin Goat Anti-Mouse at 1:2000 for 1 hour at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 200 μL for 2 minutes at RT. Magnification: 40x.



Image 2. CamKII (6G9) in cultured cortical neurons.



#### **Western Blotting**

Image 3. western rat membrane 1 in 1000 CaMKII.

Please check the product details page for more images. Overall 5 images are available for ABIN361644.