

Datasheet for ABIN361658
anti-Nitrotyrosine antibody[Go to Product page](#)

7 Images

9 Publications

Overview

Quantity:	100 µg
Target:	Nitrotyrosine
Reactivity:	Please inquire
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Nitrotyrosine antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunoprecipitation (IP), Flow Cytometry (FACS), Immunocytochemistry (ICC), Immunofluorescence (IF), Antibody Array (AA)

Product Details

Immunogen:	3-(4-hydroxy-3-nitrophenylacetamido) propionic acid-bovine serum albumin
Clone:	39B6
Isotype:	IgG2a
Specificity:	Recognizes 3-nitrotyrosine moieties. No detectable cross-reactivity with non-nitrated tyrosine. Not species specific.
Purification:	Protein G Purified

Target Details

Target:	Nitrotyrosine
Abstract:	Nitrotyrosine Products

Target Details

Target Type:	Chemical
Background:	<p>Protein tyrosine nitration results in a post-translational modification that is increasingly receiving attention as an important component of nitric oxide signaling (2). While multiple nonenzymatic mechanisms are known to be capable of producing nitrated tyrosine residues, most tyrosine nitration events involve catalysis by metalloproteins such as myeloperoxidase, eosino-philperoxidase (3), myoglobin, the cytochrome P-450s, superoxide dismutase and prostacyclin synthase. Nitrotyrosine may also serve as a biomarker for the effects of reactive nitrogen oxides, based on tyrosine residues becoming nitrated in proteins at sites of inflammation induced tissue injury (1). The presence of nitro tyrosine-containing proteins therefore has shown high correlation to disease states such as atherosclerosis, Alzheimer's disease, Parkinson's disease and amyotrophic lateral sclerosis (4).</p>

Application Details

Application Notes:	<ul style="list-style-type: none">• WB (1:1400)• IHC (1:100)• optimal dilutions for assays should be determined by the user.
Comment:	0.7 µg/ml of ABIN361657 was sufficient for detection of 5 µg SIN-1 treated BSA by Western Blot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only

Handling


Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	-20°C

Publications

Product cited in: Su, Gu, Wang, Wang: "Lidocaine attenuates proinflammatory cytokine production induced by extracellular adenosine triphosphate in cultured rat microglia." in: **Anesthesia and analgesia**, Vol. 111, Issue 3, pp. 768-74, (2010) ([PubMed](#)).

There are more publications referencing this product on: [Product page](#)

Images



Western Blotting

Image 1. Western Blot analysis of Human Recombinant Protein showing detection of Multiple Bands Nitrotyrosine protein using Mouse Anti-Nitrotyrosine Monoclonal Antibody, Clone 39B6 (ABIN361657 and ABIN361658). Lane 1: MW Ladder. Lane 2: hASYN Monomer (3.84 µg). Lane 3: Nitrosylated hASYN (3.84 µg).. Block: 5 % Skim Milk Powder in TBST. Primary Antibody: Mouse Anti-Nitrotyrosine Monoclonal Antibody (ABIN361657 and ABIN361658) at 1:1000 for 2 hours at RT with shaking. Secondary Antibody: Goat anti-mouse IgG:HRP at 1:4000 for 2 hour at RT with shaking. Color Development: Chemiluminescent for HRP (Moss) for 5 min in RT. Predicted/Observed Size: Multiple Bands.




Image 2. Nitrotyrosine (39B6), mouse backskin




Image 3. Nitrotyrosine (39B6), rat liver

Please check the [product details page](#) for more images. Overall 7 images are available for ABIN361658.