antibodies - online.com







anti-DLG4 antibody





\sim	
()\/\	rview
\circ	

Quantity:	100 μg
Target:	DLG4
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC), Antibody Array (AA)

Product Details

Immunogen:	Recombinant rat PSD-95
Clone:	7E3
Isotype:	lgG1
Specificity:	Detects \sim 100 kDa. An additional protein of >100 kDa is also detected. Additional cross-reactive bands are detected at \sim 75 kDa and 50 kDa in rat and mouse samples.
Cross-Reactivity:	Cow, Human, Mouse, Rat
Purification:	Protein G Purified

Target Details

Target:	DLG4
Alternative Name:	PSD95 (DLG4 Products)

Target Details

Concentration:


Target Details	
Background:	Postsynaptic Density protein 95 (PSD95), also known as Synapse associated protein 90 kDa, is
	a member of the membrane-associated guanylate kinase (MAGUK) family of proteins. PSD95 is
	a scaffolding protein and is involved in the assembly and function of the postsynaptic density
	complex (1). These family members consist of an N-terminal variable segment followed by
	three amino-terminal PDZ domains, an upstream SH3 domain and an inactive carboxyl-termina
	guanylate kinase (GK) domain. The first and second PDZ domain localize NMDA receptors and
	K+ channels to synapses, and the third binds to neuroligins which are neuronal cell adhesion
	molecules that interact with b-neurexins and form intercellular junctions. PSD-95 also binds to
	neuronal nitric oxide synthase, possibly through interactions between PDZ domains present on
	both proteins (2). Thus different PDZ domains of PSD-95 might be specialized for distinct
	functions (3, 4). PSD95 participates in synaptic targeting of AMPA receptors through an indirect
	manner involving Stargazin and related transmembrane AMPA receptor regulatory proteins
	(TARPs) (5). The protein is implicated in experience dependent plasticity and plays an
	indispensable role in learning (6). Mutations in PSD95 are associated with autism (7).
Gene ID:	29495
NCBI Accession:	NP_062567
UniProt:	P31016
Pathways:	Regulation of Muscle Cell Differentiation, Synaptic Membrane, Skeletal Muscle Fiber
	Development, Asymmetric Protein Localization, Regulation of long-term Neuronal Synaptic
	Plasticity
Application Details	
Application Notes:	• WB (1:1000)
	• IHC (1:1000)
	• ICC/IF (1:100)
	optimal dilutions for assays should be determined by the user.
Comment:	1 μg/ml was sufficient for detection of PSD-95 on 20 μg rat brain tissue extract by ECL
	immunoblot analysis using Goat Anti-Mouse IgG: HRP as the secondary.
Restrictions:	For Research Use only
Handling	
Format:	Liquid

1 mg/mL

Handling

Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	-20°C

Images




250 148 98 64 50 36 22 16

Immunohistochemistry

Image 1. Immunohistochemistry analysis using Mouse Anti-PSD95 Monoclonal Antibody, Clone 7E3 (ABIN361721 and ABIN361722). Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-PSD95 Monoclonal Antibody (ABIN361721 and ABIN361722) at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Basal cell staining in the epidermis, some hair follicle staining, dermal staining. Backskin obtained from transgenic mice...

Western Blotting

Image 2. western rat membrane 1 in 1000 PSD 95.



Immunohistochemistry

Image 3. Immunohistochemistry analysis using Mouse Anti-PSD95 Monoclonal Antibody, Clone 7E3 (ABIN361721 and ABIN361722). Tissue: Neocortex. Species: Rat. Primary Antibody: Mouse Anti-PSD95 Monoclonal Antibody (ABIN361721 and ABIN361722) at 1:1000.

Please check the product details page for more images. Overall 4 images are available for ABIN361722.