antibodies - online.com







anti-TRPV3 antibody (AA 774-791)



Images



| \sim | |
|-------------------------|-----------------|
| $() \setminus (\cap)$ | $r(1) \cap (1)$ |
| \cup | rview |
| | _ |

| Quantity: | 100 μg |
|----------------------|--|
| Target: | TRPV3 |
| Binding Specificity: | AA 774-791 |
| Reactivity: | Rat |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This TRPV3 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC), Immunofluorescence (IF), Antibody Array (AA) |


Product Details

| Immunogen: | Synthetic peptide amino acids 774-791 (C-terminus) of rat TrpV3 |
|-------------------|---|
| Clone: | N15-4 (Formerly S15-4) |
| Isotype: | lgG2a |
| Specificity: | Detects ~70 kDa. |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Protein G Purified |

Target Details


Target Details

| rarget Details | |
|---------------------|--|
| Alternative Name: | TRPV3 (TRPV3 Products) |
| Background: | The TRPV3 protein belongs to a family of nonselective cation channels that function in a variet of processes, including temperature sensation and vasoregulation. The thermosensitive members of this family are expressed in subsets of sensory neurons that terminate in the skin and are activated at distinct physiological temperatures. This channel is activated at temperatures between 22 and 40 degrees C. The gene lies in close proximity to another family member (TRPV1) gene on chromosome 17, and the two encoded proteins are thought to associate with each other to form heteromeric channels (1, 2). |
| Gene ID: | 497948 |
| NCBI Accession: | NP_001020928 |
| UniProt: | Q4QYD9 |
| Application Details | |
| Application Notes: | WB (1:1000) IHC (1:1000) ICC/IF (1:100) optimal dilutions for assays should be determined by the user. |
| Comment: | 1 μ g/ml of ABIN361778 was sufficient for detection of TrpV3 in 10 μ g of COS-1 cell lysate transiently transfected with TrpV3 by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 mg/mL |
| Buffer: | PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | -20°C |
| | |




Immunohistochemistry

Image 1. Immunohistochemistry analysis using Mouse Anti-TrpV3 Monoclonal Antibody, Clone N15/4 (ABIN361778). Tissue: hippocampus. Species: Human. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-TrpV3 Monoclonal Antibody (ABIN361778) at 1:1000 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT.



Immunofluorescence

Image 2. TrpV3 (S15 4), Human Hippocampus.



Immunocytochemistry

Image 3. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-TrpV3 Monoclonal Antibody, Clone N15/4 (ABIN361778). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-TrpV3 Monoclonal Antibody (ABIN361778) at 1:100 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) TrpV3 Antibody (D) Composite.

Please check the product details page for more images. Overall 5 images are available for ABIN361778.