

Datasheet for ABIN375627 **anti-Interferon gamma antibody**

1 Image



Overview

Quantity:	0.5 mg
Target:	Interferon gamma (IFNG)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Interferon gamma antibody is un-conjugated
Application:	ELISA, Flow Cytometry (FACS), Detection (D)

Product Details

Immunogen:	E. coli-expressed human IFN-
Clone:	B27
Isotype:	lgG1
Specificity:	Human/Rhesus/Cynomolgus/Chimpanzee/Pigtail Macaque/African Green Monkey/Sooty Mangabey IFN-
Characteristics:	Mouse Anti-Human IFN-g-UNLB
Purification:	Purified

Target Details

Target:	Interferon gamma (IFNG)
Alternative Name:	IFN-g (IFNG Products)

Target Details

Pathways:

Interferon-gamma Pathway, Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, ER-Nucleus Signaling, Regulation of Carbohydrate Metabolic Process, Protein targeting to Nucleus, Autophagy

Application Details

Application Notes:

- Applications: ELISA-Detection Quality tested, FC Quality tested, ELISPOT-Capture Reported in literature, IHC-FS Reported in literature, ICC Reported in literature, IP Reported in literature, Neut Reported in literature, Multiplex-Capture Reported in literature,
 Note May be paired with the purified clone A, (SB Cat. No.,) in a sandwich ELISA
- Working Dilutions: ELISA BIOT conjugate 1:2,000 1:8,000 Flow Cytometry FITC and PE conjugates 10 L/106 cells For flow cytometry, the suggested use of these reagents is in a final volume of 100 L

Sample Volume:


1 mL

Restrictions:

For Research Use only

Handling

Concentration:	0.5 mg/mL
Buffer:	0.5 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. No preservatives or amine-containing buffer salts added
Preservative:	Without preservative
Handling Advice:	Each reagent is stable for the period shown on the bottle label if stored as directed.
Storage:	4 °C
Storage Comment:	Store at 2-8°C



Flow Cytometry

Image 1. PMA and ionomycin stimulated human peripheral blood lymphocytes were stained with Mouse Anti-Human CD3-PE.