

Datasheet for ABIN375775

anti-CD3 epsilon antibody (APC)**1** Image[Go to Product page](#)

Overview

Quantity:	0.1 mg
Target:	CD3 epsilon (CD3E)
Reactivity:	Mouse
Host:	Hamster
Clonality:	Monoclonal
Conjugate:	This CD3 epsilon antibody is conjugated to APC
Application:	Flow Cytometry (FACS)

Product Details

Immunogen:	H-2Kb-specific murine cytotoxic T-lymphocyte (CTL) clone BM10-37
Clone:	145-2C11
Isotype:	IgG
Specificity:	Mouse CD3, Mr 25 kDa
Characteristics:	Hamster Anti-Mouse CD3e-APC
Purification:	Purified

Target Details

Target:	CD3 epsilon (CD3E)
Alternative Name:	CD3e (CD3E Products)
Background:	CD3 ϵ , a member of the immunoglobulin superfamily of cell surface receptors, is comprised of

Target Details

five invariable chains ranging in size from 16-28 kDa and is closely associated with the T cell antigen receptor (TCR). It is expressed on all T cells of all mouse strains. CD3 plays a major role in signaling during antigen recognition, leading to T-cell activation.

Pathways: [TCR Signaling](#), [CXCR4-mediated Signaling Events](#), [Ubiquitin Proteasome Pathway](#)

Application Details

Application Notes:

- **Applications:** FC - Quality tested , IHC - Reported in literature , IP - Reported in literature , WB - Reported in literature , Block - Reported in literature , Depletion - Reported in literature , Activ - Reported in literature , CMCD - Reported in literature
- **Working Dilutions:** Flow Cytometry FITC conjugate 3 g/106 cells BIMA and AF488 conjugates 1 g/106 cells PE, APC, SPRD, and AF647 conjugates 0.1 g/106 cells For flow cytometry, the suggested use of these reagents is in a final volume of 100 l

Comment: In vitro depletion of CD3+ cells, In vitro activation of T cells

Sample Volume: 1 mL

Restrictions: For Research Use only

Handling

Concentration: 0.1 mg/mL

Buffer: 0.1 mg in 1.0 mL of PBS/Sodium azide and a stabilizing agent


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling Advice: **Do not freeze!**
Protect conjugated products from light.
Each reagent is stable for the period shown on the bottle label if stored as directed.

Storage: 4 °C

Storage Comment: Store at 2-8°C



Flow Cytometry

Image 1. BALB/c mouse splenocytes were stained with Hamster Anti-Mouse CD3ε-BIMA.