

## Datasheet for ABIN375875

## Goat anti-Rabbit Ig (Heavy & Light Chain) Antibody (beta-Gal)






Go to Product page

| Overview             |   |
|----------------------|---|
| Quantity:            | 1 mL  |
| Target:              | Ig  |
| Binding Specificity: | Heavy & Light Chain   |
| Reactivity:          | Rabbit  |
| Host:                | Goat  |
| Clonality:           | Polyclonal  |
| Conjugate:           | beta-Gal  |
| Application:         | ELISA, Immunohistochemistry (IHC), Western Blotting (WB)  |
| Product Details      |   |
| Isotype:             | IgG   |
| Specificity:         | Reacts with the heavy and light chains of rabbit IgM and IgG, and with the light chains of rabbit IgA as demonstrated by ELISA and flow cytometry.                          |
| Characteristics:     | Source: Pooled antisera from goats hyperimmunized with normal rabbit IgM and IgG.   |
|                      | To insure lot-to-lot consistency, each batch of product is tested by ELISA, PCFIA and/or flow cytometry for conformance to characteristics of a standard reference reagent. |
| Purification:        | Affinity chromatography on human IgM covalently linked to Sepharose-4B.   |
| Target Details       |   |
| Target:              | lg  |
| Abstract:            | Ig Products   |

## **Application Details**

| Application Notes: | Each laboratory should determine an optimum working titer for use in its particular application.  Other applications have not been tested but use in such assays should not necessarily be excluded. |
|--------------------|--|
| Restrictions:      | For Research Use only  |
| Handling           |  |
| Format:            | Liquid   |
| Handling Advice:   | A vaid was a stad for a min or and the avvisor   |
| Harlaning Advice.  | Avoid repeated freezing and thawing.   |
| Storage:           | 4 °C   |



**Image 1.** Lysates from mouse embryonic fibroblasts expressing no Bak (Bax-/-Bak-/- (DKO)), mouse Bak (Bax-/-), or WT human Bak (in DKO) were resolved by electrophoresis, transferred to nitrocellulose membrane, and probed with anti-Bak followed by Goat Anti-Rabbit Ig, Human ads-HRP