

Datasheet for ABIN375963

Goat anti-Human lambda (Chain lambda) Antibody (Biotin)





Go to Product page

Overview	
Quantity:	0.5 mg
Target:	lambda
Binding Specificity:	Chain lambda
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	Biotin
Application:	Flow Cytometry (FACS), ELISA
Product Details	
Isotype:	IgG
Fragment:	F(ab')2 fragment
Specificity:	Reacts with humank light chains as demonstrated by ELISA and flow cytometry.
Cross-Reactivity (Details):	Cross Absorption: Pooled human myeloma proteins withk light chains and mouse immunoglobulins.
Characteristics:	Goat F(ab')2 Anti-Human Lambda-BIOT
Purification:	Affinity chromatography on pooled human Igs with λ light chains covalently linked to agarose.
Target Details	
Target:	lambda

Application Details

- · Applications: Quality tested applications include ELISA FLISA FC,
- · Other referenced applications include WB, Stim
- Working Dilutions: ELISA HRP conjugate 1:4,000 1:8,000 BIOT conjugate 1:5,000 1:20,000 FLISA FITC conjugate 1:200 1:400 PE and AF647 conjugates 1 g/mL Flow Cytometry FITC and BIOT conjugates 1 g/106 cells PE and AF647 conjugates 0.1 g/106 cells For flow cytometry, the suggested use of these reagents is in a final volume of 100 L

Comment: B cell enumeration


Sample Volume: 1 mL

Restrictions: For Research Use only

Handling

Concentration:	0.5 mg/mL
Buffer:	0.5 mg in 1.0 mL of PBS/Sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect conjugated products from light. Each reagent is stable for the period shown on the bottle label if stored as directed.
Storage:	4 °C
Storage Comment:	Store at 2-8°C

Images



ELISA

Image 1. ELISA plate was coated with purified human IgG κ , IgM κ , IgA κ , IgG λ , IgM λ , and IgA λ . Immunoglobulins were detected with serially diluted Goat F(ab')2 Anti-Human Lambda-BIOT.