

Datasheet for ABIN376277

Goat anti-Rat Ig (Heavy & Light Chain) Antibody (HRP)





Go to Product page

Overview	
Quantity:	1 mL
Target:	lg
Binding Specificity:	Heavy & Light Chain
Reactivity:	Rat
Host:	Goat
Clonality:	Polyclonal
Conjugate:	HRP
Application:	ELISA, Immunohistochemistry (IHC), Western Blotting (WB)
Product Details	
Isotype:	IgG
Specificity:	Reacts with the heavy and light chains of rat IgM and IgG, and with the light chains of rat IgA as
	demonstrated by ELISA and/or flow cytometry.
Characteristics:	demonstrated by ELISA and/or flow cytometry. Source: Pooled antisera from goats hyperimmunized with normal rat IgM and IgG
Characteristics:	
Characteristics:	Source: Pooled antisera from goats hyperimmunized with normal rat IgM and IgG
Characteristics: Purification:	Source: Pooled antisera from goats hyperimmunized with normal rat IgM and IgG To insure lot-to-lot consistency, each batch of product is tested by ELISA, PCFIA and/or flow
	Source: Pooled antisera from goats hyperimmunized with normal rat IgM and IgG To insure lot-to-lot consistency, each batch of product is tested by ELISA, PCFIA and/or flow cytometry for conformance to characteristics of a standard reference reagent.
Purification:	Source: Pooled antisera from goats hyperimmunized with normal rat IgM and IgG To insure lot-to-lot consistency, each batch of product is tested by ELISA, PCFIA and/or flow cytometry for conformance to characteristics of a standard reference reagent.

Application Details

Application Notes:	Working Dilution:
	1:4,000-1:8,000
	Representative data are included in this product insert.
	Each laboratory should determine an optimum working titer for use in its particular application.
	Other applications have not been tested but use in such assays should not necessarily be
	excluded.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Handling Advice:	Avoid repeated freezing and thawing. Dilute only prior to immediate use
	Centrifuge product if not completely clear after standing at room temperature.
	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish
	peroxidase.
Storage:	4 °C




Image 1. Lysates from mouse embryonic fibroblasts expressing no Bak (Bax-/-Bak-/- (DKO)), mouse Bak (Bax-/-), or WT human Bak (in DKO) were resolved by electrophoresis, transferred to nitrocellulose membrane, and probed with anti-Bak followed by Goat Anti-Rat Ig, Mouse ads-HRP