

Datasheet for ABIN376287

**Goat anti-Rabbit IgM (Heavy Chain) Antibody (HRP)**[Go to Product page](#)**1** Image

## Overview

|                      |  |
|----------------------|--|
| Quantity:            | 1 mL   |
| Target:              | IgM  |
| Binding Specificity: | Heavy Chain  |
| Reactivity:          | Rabbit   |
| Host:                | Goat   |
| Clonality:           | Polyclonal   |
| Conjugate:           | HRP  |
| Application:         | ELISA, Immunohistochemistry (IHC), Western Blotting (WB) |

## Product Details

|                  |   |
|------------------|---|
| Isotype:         | IgG   |
| Specificity:     | Reacts with the heavy chain of rabbit IgM as demonstrated by ELISA.   |
| Characteristics: | Source: Pooled antisera from goats hyperimmunized with normal rabbit IgM<br>To insure lot-to-lot consistency, each batch of product is tested by ELISA, PCFIA and/or flow cytometry for conformance to characteristics of a standard reference reagent. |
| Purification:    | Affinity chromatography on rabbit IgG covalently linked to agarose  |

## Target Details

|           |                              |
|-----------|------------------------------|
| Target:   | IgM                          |
| Abstract: | <a href="#">IgM Products</a> |

## Target Details

Target Type: Antibody

## Application Details

Application Notes: Working Dilution:  
1:4,000-1:8,000  
Representative data are included in this product insert.

Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded.

Restrictions: For Research Use only


## Handling

Format: Liquid

Handling Advice: Avoid repeated freezing and thawing. Dilute only prior to immediate use  
Centrifuge product if not completely clear after standing at room temperature.  
Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.

Storage: 4 °C

## Images



### ELISA

**Image 1.** ELISA plate was coated with purified rabbit IgM and IgG. Immunoglobulins were detected with serially diluted Goat Anti-Rabbit IgM-HRP.