

## Datasheet for ABIN376302

## Rabbit anti-Goat IgG (Heavy & Light Chain) Antibody (HRP)



Go to Product page

| Overview             |  |
|----------------------|--|
| Quantity:            | 1 mL   |
| Target:              | IgG  |
| Binding Specificity: | Heavy & Light Chain  |
| Reactivity:          | Goat   |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | HRP  |
| Application:         | ELISA, Immunohistochemistry (IHC), Western Blotting (WB)   |
| Product Details      |  |
| Isotype:             | IgG  |
| Fragment:            | F(ab')2 fragment   |
| Specificity:         | Reacts with the heavy and light chains of goat IgG as demonstrated by ELISA.   |
| Characteristics:     | Source: Pepsin digest of rabbit anti-goat IgG (H+L).  To insure lot-to-lot consistency, each batch of product is tested by ELISA and PCFIA for conformance to characteristics of a standard reference reagent. |
| Purification:        | Affinity chromatography on hamster IgG covalently linked to agarose. F(ab') 2 fragment prepared by pepsin digestion of affinity purified IgG.  |
| Target Details       |  |
| Target:              | IgG  |

## **Target Details**

| Abstract:           | IgG Products   |
|---------------------|--|
| Target Type:        | Antibody   |
|                     |  |
| Application Details |  |
| Application Notes:  | Working Dilution:  |
|                     | 1:4,000-1:8,000  |
|                     | Representative data are included in this product insert.   |
|                     | Each laboratory should determine an optimum working titer for use in its particular application. |
|                     | Other applications have not been tested but use in such assays should not necessarily be         |
|                     | excluded.  |
| Restrictions:       | For Research Use only  |
| Handling            |  |
| Format:             | Liquid   |
| Handling Advice:    | Avoid repeated freezing and thawing. Dilute only prior to immediate use                          |
|                     | Centrifuge product if not completely clear after standing at room temperature.                   |
|                     | Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish         |
|                     | peroxidase.  |
| Storage:            | 4 °C   |