



[Go to Product page](#)

Datasheet for ABIN376754

## Goat anti-Mouse IgG2b (Heavy Chain) Antibody (TRITC)

1 Image

6 Publications

### Overview

Quantity:	1 mg
Target:	IgG2b
Binding Specificity:	Heavy Chain
Reactivity:	Mouse
Host:	Goat
Clonality:	Polyclonal
Conjugate:	TRITC
Application:	ELISA, Immunohistochemistry (IHC), Flow Cytometry (FACS)

### Product Details

Isotype:	IgG
Specificity:	Reacts with the heavy chain of mouse IgG 2b as demonstrated by ELISA and flow cytometry.
Characteristics:	Source: Pooled antisera from goats hyperimmunized with mouse IgG 2b paraproteins. To insure lot-to-lot consistency, each batch of product is tested by ELISA, PCFIA and/or flow cytometry for conformance to characteristics of a standard reference reagent.
Purification:	Affinity chromatography on mouse IgG 1 covalently linked to agarose.

### Target Details

Target:	IgG2b
Abstract:	<a href="#">IgG2b Products</a>

## Target Details

---

Target Type: Antibody

## Application Details

---

Application Notes: Working Dilution:  
<= 1 µg/10<sup>6</sup> cells  
Representative data are included in this product insert.

Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded.

Restrictions: For Research Use only

## Handling

---

Handling Advice: **Protect conjugated products from light.**  
Each reagent is stable for the period shown on the bottle label if stored as directed.


Storage: 4 °C

## Publications

---

Product cited in: Pietrek, Brinkmann, Glowacka, Enlund, Haevemeier, Dittrich-Breiholz, Kracht, Lewitzky, Saksela, Feller, Schulz: "The role of the Kaposi Sarcoma Herpesvirus K15 SH3 binding site in inflammatory signaling and B-cell activation." in: **Journal of virology**, (2010) ([PubMed](#)).

There are more publications referencing this product on: [Product page](#)



### Immunohistochemistry

**Image 1.** Paraffin embedded normal breast tissue section was stained with anti-acetylated tubulin (IgG2b) and  $\gamma$ -tubulin (IgG1) followed by Goat Anti-Mouse IgG2b, Human ads-TRITC