antibodies -online.com





Goat anti-Rat IgG (Heavy Chain) Antibody (TRITC)





Go to Product page

| Overview | |
|----------------------|---|
| Quantity: | 1 mg |
| Target: | IgG |
| Binding Specificity: | Heavy Chain |
| Reactivity: | Rat |
| Host: | Goat |
| Clonality: | Polyclonal |
| Conjugate: | TRITC |
| Application: | ELISA, Immunohistochemistry (IHC), Flow Cytometry (FACS) |
| Product Details | |
| Isotype: | IgG |
| Specificity: | Reacts with the heavy chain of rat IgG as demonstrated by ELISA. |
| Characteristics: | Source: Pooled antisera from goats hyperimmunized with normal rat IgG |
| | To insure lot-to-lot consistency, each batch of product is tested by ELISA, PCFIA and/or flow |
| | cytometry for conformance to characteristics of a standard reference reagent. |
| Purification: | Affinity chromatography on sheep IgG covalently linked to agarose. |
| Target Details | |
| Target: | IgG |
| Abstract: | IgG Products |
| | |

Target Details

Target Type:

Antibody

Application Details

Application Notes:

Working Dilution:

<= 1 µg/10^6 cells

Representative data are included in this product insert.

Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded.

Restrictions:

For Research Use only

Handling

Handling Advice:


Protect conjugated products from light.

Each reagent is stable for the period shown on the bottle label if stored as directed.

Storage:

4°C

Images



ELISA

Image 1. FLISA plate was coated with purified rat IgG and IgM. Immunoglobulins were detected with serially diluted Goat Anti-Rat IgG-TRITC.