antibodies -online.com





Goat anti-Rabbit IgG (Heavy & Light Chain) Antibody (TRITC)



1 Image	2 Publications
Overview	
Quantity:	1 mg

IgG

Heavy & Light Chain

Reactivity:	Rabbit

Host:	Goat

Clonality:	Polyclonal
------------	------------

Conjugate:	TRITC
------------	-------

Application:	ELISA, Immunohistochemistry	/ (IHC), Flow C	ytometry ((FACS)	

Product Details

Target:

Binding Specificity:

Isotype:	IgG
Specificity:	Reacts with the heavy and light chains of rabbit IgG, and with the light chains of rabbit IgM and IgA as demonstrated by ELISA and flow cytometry.
Characteristics:	Source: Pooled antisera from goats hyperimmunized with normal rabbit IgG. To insure lot-to-lot consistency, each batch of product is tested by ELISA, PCFIA and/or flow cytometry for conformance to characteristics of a standard reference reagent.
Purification:	Affinity chromatography on pooled human IgG covalently linked to agarose.

Target Details

Target:	IgG
Abstract:	IgG Products

Target Details Antibody Target Type: **Application Details** Application Notes: Working Dilution: <= 1 μg/10⁶ cells Representative data are included in this product insert. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded. Restrictions: For Research Use only Handling Handling Advice: Protect conjugated products from light. Each reagent is stable for the period shown on the bottle label if stored as directed. Storage: 4°C **Publications**

Product cited in:

Skowron-Kandzia, Tomsia, Koryciak-Komarska, Plewka, Wieczorek, Czekaj: "Gene Expression in Amnion-Derived Cells Cultured on Recombinant Laminin 332-A Preliminary Study." in: **Frontiers in medicine**, Vol. 8, pp. 719899, (2021) (PubMed).


Leiendecker, Jung, Krecioch, Neumann, Schleiffer, Mechtler, Wiesner, Obenauf: "LSD1 inhibition induces differentiation and cell death in Merkel cell carcinoma." in: **EMBO molecular medicine**, pp. e12525, (2020) (PubMed).

Shan, Xu, Bleyer, Becker, Melbaum, Wemheuer, Hirschfeld, Wacker, Zhao, Schütz, Brenig: " Association of α/β -Hydrolase D16B with Bovine Conception Rate and Sperm Plasma Membrane Lipid Composition." in: **International journal of molecular sciences**, Vol. 21, Issue 2, (2020) (PubMed).

Frenzel, Willbold: "Kinetic titration series with biolayer interferometry." in: **PLoS ONE**, Vol. 9, Issue 9, pp. e106882, (2016) (PubMed).

Aeckerle, Drummer, Debowski, Viebahn, Behr: "Primordial germ cell development in the marmoset monkey as revealed by pluripotency factor expression: suggestion of a novel model of embryonic germ cell translocation." in: **Molecular human reproduction**, Vol. 21, Issue 1, pp. 66-80, (2015) (PubMed).

Images



Immunohistochemistry

Image 1. Paraffin embedded human kidney section from AAV patient was stained with anti-C5L2 followed by Goat Anti-Rabbit IgG(H+L), Mouse/Human ads-TRITC