antibodies .- online.com







Goat IgG Isotype Control

Image

Publications



Overview

Quantity:	10 mg
Target:	IgG
Host:	Goat
Clonality:	Polyclonal
Application:	ELISA, Flow Cytometry (FACS)

Product Details

1 Toddot Details		
Isotype:	IgG	
Characteristics:	Goat IgG-UNLB	
Purification:	Purified	
Target Details		
Target:	IgG	
Abstract:	IgG Products	
Target Type:	Antibody	

Application Details


Application Notes:

- · Applications: Quality tested applications include FC, ELISA FLISA
- Other referenced applications include IHC-FS , IHC-PS , IHC-WM , ICC , IP , Block , In vitro control, In vivo control
- Working Dilutions: Flow Cytometry BIOT conjugate 1 g/106 cells FITC, PE, APC, SPRD, CY5,

Application Details

Application Details	
	and PE/CY5.5 conjugates 10 L/106 cells For flow cytometry, the suggested use of these reagents is in a final volume of 100 L FLISA TRITC and TXRD conjugates 1:100 - 1:400
Sample Volume:	2 mL
Restrictions:	For Research Use only
Handling	
Concentration:	5 mg/mL
Buffer:	10.0 mg of purified immunoglobulin in 2.0 mL of borate buffered saline, pH 8.2. No preservatives or amine-containing buffer salts added
Preservative:	Without preservative
Handling Advice:	Each reagent is stable for the period shown on the bottle label if stored as directed.
Storage:	4 °C
Storage Comment:	Store at 2-8°C
Publications	
Product cited in:	Aeckerle, Drummer, Debowski, Viebahn, Behr: "Primordial germ cell development in the
	marmoset monkey as revealed by pluripotency factor expression: suggestion of a novel model
	of embryonic germ cell translocation." in: Molecular human reproduction , Vol. 21, Issue 1, pp.
	66-80, (2015) (PubMed).
	Saiya-Cork, Collins, Parkin, Ouillette, Kuizon, Kujawski, Erba, Campagnaro, Shedden, Kaminski,
	Malek: "A pathobiological role of the insulin receptor in chronic lymphocytic leukemia." in:
	Clinical cancer research : an official journal of the American Association for Cancer

Research, Vol. 17, Issue 9, pp. 2679-92, (2011) (PubMed).



Immunohistochemistry

Image 1. Paraffin embedded human gastric cancer tissue was stained with Goat IgG-UNLB isotype control, DAB, and hematoxylin.