

Datasheet for ABIN376897

Goat anti-Human IgD Antibody





Overview

Overview	
Quantity:	0.5 mg
Target:	IgD
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Application:	Flow Cytometry (FACS), ELISA
Product Details	
Isotype:	IgG
Fragment:	F(ab')2 fragment
Specificity:	Reacts with the heavy chain of human IgD
Characteristics:	Goat F(ab')2 Anti-Human IgD-UNLB
Purification:	Purified
Target Details	
Target:	IgD
Abstract:	IgD Products
Target Type:	Antibody

Application Details

Application Notes:

- Applications: Quality tested applications include ELISA FLISA FC,
- Other referenced applications include IHC-FS , IHC-PS , Sep , Stim
- Working Dilutions: ELISA BIOT conjugate 1:5,000 1:20,000 FLISA FITC, AF488, and AF555 conjugates 1:100 1:400 PE and AF647 conjugates 1 g/mL Flow Cytometry FITC, BIOT, and AF488 conjugates 1 g/106 cells PE and AF647 conjugates 0.1 g/106 cells For flow cytometry, the suggested use of these reagents is in a final volume of 100 L


Restrictions:

For Research Use only

Handling

Concentration:	0.5 mg/mL
Buffer:	0.5 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. No preservatives or amine-containing buffer salts added
Preservative:	Without preservative
Handling Advice:	Each reagent is stable for the period shown on the bottle label if stored as directed.
Storage:	4 °C
Storage Comment:	Store at 2-8°C

Images



ELISA

Image 1. FLISA plate was coated with purified human IgD, IgG, IgM, and IgA. Immunoglobulins were detected with serially diluted Goat F(ab')2 Anti-Human IgD-AF488.