

Datasheet for ABIN376914

Goat anti-Rat Ig (Heavy & Light Chain) Antibody






Go to Product page

Overview	
Quantity:	1 mg
Target:	lg
Binding Specificity:	Heavy & Light Chain
Reactivity:	Rat
Host:	Goat
Clonality:	Polyclonal
Application:	ELISA, Flow Cytometry (FACS), Immunohistochemistry (IHC), Western Blotting (WB)
Product Details	
Isotype:	IgG
Isotype: Specificity:	IgG Reacts with the heavy and light chains of rat IgM and IgG, and with the light chains of rat IgA as demonstrated by ELISA and/or flow cytometry.
	Reacts with the heavy and light chains of rat IgM and IgG, and with the light chains of rat IgA as
Specificity:	Reacts with the heavy and light chains of rat IgM and IgG, and with the light chains of rat IgA as demonstrated by ELISA and/or flow cytometry. Source: Pooled antisera from goats hyperimmunized with normal rat IgM and IgG To insure lot-to-lot consistency, each batch of product is tested by ELISA, PCFIA and/or flow
Specificity: Characteristics:	Reacts with the heavy and light chains of rat IgM and IgG, and with the light chains of rat IgA as demonstrated by ELISA and/or flow cytometry. Source: Pooled antisera from goats hyperimmunized with normal rat IgM and IgG To insure lot-to-lot consistency, each batch of product is tested by ELISA, PCFIA and/or flow cytometry for conformance to characteristics of a standard reference reagent.
Specificity: Characteristics: Purification:	Reacts with the heavy and light chains of rat IgM and IgG, and with the light chains of rat IgA as demonstrated by ELISA and/or flow cytometry. Source: Pooled antisera from goats hyperimmunized with normal rat IgM and IgG To insure lot-to-lot consistency, each batch of product is tested by ELISA, PCFIA and/or flow cytometry for conformance to characteristics of a standard reference reagent.

Application Details

Application Notes:	Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded.
Restrictions:	For Research Use only
Handling	
Buffer:	1.0 mg purified immunoglobulin in 1.0 mL of 100 mM borate buffered saline, pH 8.2.
Preservative:	Azide free
Handling Advice:	Each reagent is stable for the period shown on the bottle label if stored as directed.
Storage:	4 °C

Images



Flow Cytometry

Image 1. BALB/c mouse splenocytes were stained with Rat Anti-Mouse CD45-UNLB.