

Datasheet for ABIN376927

Goat anti-Rabbit Ig (Heavy & Light Chain) Antibody





Go to Product page

Overview	
Quantity:	1 mg
Target:	lg
Binding Specificity:	Heavy & Light Chain
Reactivity:	Rabbit
Host:	Goat
Clonality:	Polyclonal
Application:	ELISA, Flow Cytometry (FACS), Immunohistochemistry (IHC), Western Blotting (WB)
Product Details	
Isotype:	IgG
Specificity:	Reacts with the heavy and light chains of rabbit IgM and IgG, and with the light chains of rabbit
	IgA as demonstrated by ELISA and flow cytometry.
Characteristics:	IgA as demonstrated by ELISA and flow cytometry. Source: Pooled antisera from goats hyperimmunized with normal rabbit IgM and IgG. To insure lot-to-lot consistency, each batch of product is tested by ELISA, PCFIA and/or flow cytometry for conformance to characteristics of a standard reference reagent.
Characteristics: Purification:	Source: Pooled antisera from goats hyperimmunized with normal rabbit IgM and IgG. To insure lot-to-lot consistency, each batch of product is tested by ELISA, PCFIA and/or flow
	Source: Pooled antisera from goats hyperimmunized with normal rabbit IgM and IgG. To insure lot-to-lot consistency, each batch of product is tested by ELISA, PCFIA and/or flow cytometry for conformance to characteristics of a standard reference reagent.
Purification:	Source: Pooled antisera from goats hyperimmunized with normal rabbit IgM and IgG. To insure lot-to-lot consistency, each batch of product is tested by ELISA, PCFIA and/or flow cytometry for conformance to characteristics of a standard reference reagent.

Application Details

Application Notes:	Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded.
Restrictions:	For Research Use only
Handling	
Buffer:	1.0 mg purified immunoglobulin in 1.0 mL of 100 mM borate buffered saline, pH 8.2.
Preservative:	Azide free
Handling Advice:	Each reagent is stable for the period shown on the bottle label if stored as directed.
Storage:	4 °C
Images	




Image 1. Lysates from mouse embryonic fibroblasts expressing no Bak (Bax-/-Bak-/- (DKO)), mouse Bak (Bax-/-), or WT human Bak (in DKO) were resolved by electrophoresis, transferred to nitrocellulose membrane, and probed with anti-Bak followed by Goat Anti-Rabbit Ig, Human ads-HRP