## antibodies .- online.com







## Donkey anti-Goat IgG (Heavy & Light Chain) Antibody






| Overview   |  |
|--|--|
| Quantity:  | 1 mg   |
| Target:  | IgG  |
| Binding Specificity:   | Heavy & Light Chain  |
| Reactivity:  | Goat   |
| Host:  | Donkey   |
| Clonality:   | Polyclonal   |
| Application:   | ELISA, Immunohistochemistry (IHC), Western Blotting (WB), Flow Cytometry (FACS)  |
| Product Details  |  |
|  |  |
| Isotype:   | IgG  |
| Isotype: Specificity:  | Reacts with the heavy and light chains of Goat IgG as demonstrated by ELISA.   |
|  |  |
| Specificity:   | Reacts with the heavy and light chains of Goat IgG as demonstrated by ELISA.  Source: Pooled antisera from donkey hyperimmunized with goat IgG.  To insure lot-to-lot consistency, each batch of product is tested by ELISA and/or PCFIA for   |
| Specificity: Characteristics:                                | Reacts with the heavy and light chains of Goat IgG as demonstrated by ELISA.  Source: Pooled antisera from donkey hyperimmunized with goat IgG.  To insure lot-to-lot consistency, each batch of product is tested by ELISA and/or PCFIA for conformance to characteristics of a standard reference reagent.           |
| Specificity: Characteristics: Purification:                  | Reacts with the heavy and light chains of Goat IgG as demonstrated by ELISA.  Source: Pooled antisera from donkey hyperimmunized with goat IgG.  To insure lot-to-lot consistency, each batch of product is tested by ELISA and/or PCFIA for conformance to characteristics of a standard reference reagent.           |
| Specificity: Characteristics:  Purification:  Target Details | Reacts with the heavy and light chains of Goat IgG as demonstrated by ELISA.  Source: Pooled antisera from donkey hyperimmunized with goat IgG.  To insure lot-to-lot consistency, each batch of product is tested by ELISA and/or PCFIA for conformance to characteristics of a standard reference reagent.  Purified |

## **Application Details**

| Application Notes: | Each laboratory should determine an optimum working titer for use in its particular application.  Other applications have not been tested but use in such assays should not necessarily be excluded. |
|--------------------|--|
| Restrictions:      | For Research Use only  |
| Handling           |  |
| Buffer:            | 1.0 mg purified immunoglobulin in 1.0 mL of 100 mM borate buffered saline, pH 8.2.   |
| Preservative:      | Azide free   |
| Handling Advice:   | Each reagent is stable for the period shown on the bottle label if stored as directed.   |
| Storage:           | 4 °C   |
| Images             |  |



## Immunofluorescence

**Image 1.** Frozen newborn mouse cartilage section was stained with Goat Anti-Type II Collagen-UNLB and DAPI.