

Datasheet for ABIN387871
anti-JMJD2D antibody (C-Term)[Go to Product page](#)

1 Image

Overview

Quantity:	400 µL
Target:	JMJD2D (KDM4D)
Binding Specificity:	AA 484-516, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This JMJD2D antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	This JMJD2D antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 484-516 amino acids from the C-terminal region of human JMJD2D.
Clone:	RB10797
Isotype:	Ig Fraction
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Target Details

Target:	JMJD2D (KDM4D)
Alternative Name:	JMJD2D (KDM4D Products)

Target Details


Background:	Covalent modification of histones plays critical role in regulating chromatin structure and transcription. While most covalent histone modifications are reversible, only recently has it been established that methyl groups are subject to enzymatic removal from histones. A family of novel JmjC domain-containing histone demethylation (JHDM) enzymes have been identified that perform this specific function. Histone demethylation by JHDM proteins requires cofactors Fe(II) and alpha-ketoglutarate. Family members include JHDM1 (demethylating histone 3 at lysine 36), and JHDM2A as well as JMJD2CH3K9 (both of which demethylate histone 3 at lysine 9). Contributions of histone demethylase activity to tumor development, decreases in cell proliferation, and hormone-dependent transcriptional activation have been observed.
Molecular Weight:	58603
Gene ID:	55693
NCBI Accession:	NP_060509
UniProt:	Q6B0I6
Pathways:	Warburg Effect

Application Details

Application Notes:	WB: 1:1000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months



Western Blotting

Image 1. Western blot analysis of anti-JMJD2D (C-term) Pab in HeLa cell line lysate. JMJD2D (C-term)(arrow) was detected using the purified Pab.