antibodies -online.com







anti-CBLC antibody (C-Term)

Images



Overview

0.0	
Quantity:	400 μL
Target:	CBLC
Binding Specificity:	AA 424-453, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CBLC antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This CBLC antibody is generated from rabbits immunized with a KLH conjugated synthetic

Immunogen:	This CBLC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 424-453 amino acids from the C-terminal region of human CBLC.
Clone:	RB0696
Isotype:	lg Fraction
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Target Details


Target:	CBLC
Alternative Name:	CBLC (CBLC Products)


Target Details

Expiry Date:

6 months

rarget betails	
Background:	Cbl proteins are a family of ubiquitin protein ligases (E3s) that negatively regulate signaling by targeting activated tyrosine kinases for degradation. Cbl- c is the most recently cloned member of the Cbl proteins and is expressed only in epithelial cells (the other Cbl proteins are ubiquitously expressed). Cbl-c, like the other mammalian Cbl proteins, can ubiquitinate the activated EGFR and target it for degradation. Through interactions with proteins containing SRC homology-2 (SH2) and SH3 domains, CBL proteins modulate downstream cell signaling.
Molecular Weight:	52456
Gene ID:	23624
NCBI Accession:	NP_001124324, NP_036248
UniProt:	Q9ULV8
Pathways:	EGFR Signaling Pathway
Application Details	
Application Notes:	WB: 1:1000. IHC-P: 1:50~100
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.





Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated. BC = breast carcinoma, HC = hepatocarcinoma.

Western Blotting

Image 2. CBLC Antibody (ABIN388072 and ABIN2845789) western blot analysis in K562 cell line lysates (35 μ g/lane).This demonstrates the CBLC antibody detected the CBLC protein (arrow).