# antibodies - online.com







## anti-IL29 antibody (AA 54-85)






Overview	
Quantity:	400 μL
Target:	IL29
Binding Specificity:	AA 54-85
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IL29 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This IL29 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 54-85 amino acids from the Central region of human IL29.
Clone:	RB03233
Isotype:	Ig Fraction
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Target Details	
Target:	IL29
Alternative Name:	IL29 (IL29 Products)

## **Target Details**

Expiry Date:

6 months

9	
Background:	IL29 is a cytokine distantly related to type I interferons and the IL-10 family. This gene, interleukin 28A (IL28A), and interleukin 28B (IL28B) are three closely related cytokine genes that form a cytokine gene cluster on a chromosomal region mapped to 19q13. Expression of the cytokines encoded by the three genes can be induced by viral infection. All three cytokines have been shown to interact with a heterodimeric class II cytokine receptor that consists of interleukin 10 receptor, beta (IL10RB) and interleukin 28 receptor, alpha (IL28RA).
Molecular Weight:	21898
Gene ID:	282618
NCBI Accession:	NP_742152
UniProt:	Q8IU54
Application Details	
Application Notes:	WB: 1:1000. IHC-P: 1:50~100
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.



### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated. BC = breast carcinoma, HC = hepatocarcinoma.

#### **Western Blotting**

**Image 2.** The anti-IL29 Pab (ABIN388447 and ABIN2848744) is used in Western blot to detect IL29 in Jurkat cell lysate.