antibodies - online.com







anti-ATG4C antibody (N-Term)





\sim	
()\/\	rview
\cup	

Overview	
Quantity:	400 μL
Target:	ATG4C
Binding Specificity:	AA 33-62, N-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ATG4C antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This ATG4C antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 33-62 amino acids from the N-terminal region of human ATG4C.
Clone:	RB7561
Isotype:	lg Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target Details	
Target:	ATG4C
Alternative Name:	ATG4C (ATG4C Products)
Background:	Macroautophagy is the major inducible pathway for the general turnover of cytoplasmic

constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic enzymes and organelles during nutrient starvation. Macroautophagy involves the formation of double-membrane bound autophagosomes which enclose the cytoplasmic constituent targeted for degradation in a membrane bound structure, which then fuse with the lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded within the lysosome (or vacuole). APG4 is a cysteine protease required for autophagy, which cleaves the C-terminal part of either MAP1LC3, GABARAPL2 or GABARAP, allowing the liberation of form I. A subpopulation of form I is subsequently converted to a smaller form (form II). Form II, with a revealed C-terminal glycine, is considered to be the phosphatidylethanolamine (PE)-conjugated form, and has the capacity for the binding to autophagosomes.


Molecular Weight:	52497
Gene ID:	84938
NCBI Accession:	NP_116241, NP_835739
UniProt:	Q96DT6
Pathways:	Autophagy

Application Details


Application Notes:	WB: 1:1000. WB: 1:1000. WB: 1:1000. IHC-P: 1:50~100
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months



1 2 130 72 55 ••• 36 28



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated. BC = breast carcinoma, HC = hepatocarcinoma.

Western Blotting

Image 2. Western blot analysis of anti-hG4C-Y48 Pab 1810a in 293 cell line lysates transiently transfected with the ATG4C gene ($2 \mu g$ /lane). hG4C-Y48(arrow) was detected using the purified Pab.

Western Blotting

Image 3. The anti-G4C Pab 1810a is used in Western blot to detect G4C in mouse liver tissue lysate

Please check the product details page for more images. Overall 4 images are available for ABIN388503.