# antibodies -online.com





# anti-ZNF395 antibody (AA 141-170)

2 Images



$\sim$				٠			
	۱۱/	Δ	rv	I		1/	1
$\cup$	v	C	ΙV	ı	$\overline{}$	٧	V

Overview		
Quantity:	400 μL	
Target:	ZNF395	
Binding Specificity:	AA 141-170	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ZNF395 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	
Product Details		
Immunogen:	This HDBP2(HDRF-2) antibody is generated from rabbits immunized with a KLH conjugated	
	synthetic peptide between 141-170 amino acids from the Central region of human	
	HDBP2(HDRF-2).	
Clone:	RB9478	
Isotype:	lg Fraction	
Purification:	This antibody is purified through a protein G column, followed by dialysis against PBS.	
Target Details		
- Target Details		
Target:	ZNF395	
Alternative Name:	HDBP2 (HDRF-2) (ZNF395 Products)	

## **Target Details**


Background:	Papillomavirus Regulatory Factor 1/HDBP2 is a novel transcription factor shuttling between nucleus and cytoplasm and binds to the specific GCCGGCG, which is an essential cis-element for Huntington's disease gene expression.	
Gene ID:	55893	
NCBI Accession:	NP_061130	
UniProt:	Q9H8N7	

## Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:50~100
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Buffer:	Purified monoclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months



#### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** Formalin-fixed and paraffin-embedded human Hepatocellular carcinoma reacted with anti-HDBP2(HDRF-2) Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated.



#### **Western Blotting**

**Image 2.** Western blot analysis of anti-HDBP2(HDRF-2) Antibody (Center) (ABIN388701 and ABIN2838790) in Y79 cell line lysates (35  $\mu$ g/lane). HDBP2(arrow) was detected using the purified Pab.