antibodies - online.com





anti-ube3a antibody (C-Term)





Publication



0	1 /	-	K	/1	-	1 A
u	\/	\vdash	I \	/ I	\vdash	1/1

Quantity:	400 μL
Target:	ube3a
Binding Specificity:	AA 836-865, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ube3a antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Flow Cytometry (FACS)
Product Details	

Immunogen:	This UBE3A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 836-865 amino acids from the C-terminal region of human UBE3A.
Clone:	RB4440
Isotype:	lg Fraction
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Target Details

Target:	ube3a
Alternative Name:	UBE3A (ube3a Products)

Background:

UBE3A interacts with the E6 protein of the cancer-associated human papillomavirus types 16 and 18. The E6/E6-AP complex binds to and targets the p53 tumor-suppressor protein for ubiquitin-mediated proteolysis. It is an E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. It can target itself for ubiquitination in vitro and efficiently promotes its own degradation in vivo. It appears that only unmodified E6-AP molecules can bind efficiently to p53 in the presence of the HPV E6 oncoprotein. UBE3A binds UBQLN1 and UBQLN2. Defects in UBE3A are a cause of Angelman syndrome (AS) [MIM:105830], also known as 'happy puppet syndrome'. AS is characterized by features of severe motor and intellectual retardation, microcephaly, ataxia, frequent jerky limb movements and flapping of the arms and hands, hypotonia, hyperactivity, hypopigmentation, seizures, absence of speech, frequent smiling and episodes of paroxysmal laughter, and an unusual facies characterized by macrostomia, a large mandible and open-mouthed expression, a great propensity for protruding the tongue ('tongue thrusting'), and an occipital groove. UBE3A contains 1 HECT-type E3 ubiquitin-protein ligase domain.

Molecular Weight:	100688
Gene ID:	7337
NCBI Accession:	NP_000453, NP_570853, NP_570854
UniProt:	Q05086
Pathways:	Intracellular Steroid Hormone Receptor Signaling Pathway

Application Details

Application Notes:	IF: 1:10~50. WB: 1:1000. IHC-P: 1:50~100. FC: 1:10~50
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling

Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months


Publications

Product cited in:

Wu, Li, Fang, Yi, Chen, Long, Gao, Wei, Chen: "Investigation of synergistic mechanism and identification of interaction site of aldose reductase with the combination of gigantol and syringic acid for prevention of diabetic cataract." in: **BMC complementary and alternative medicine**, Vol. 16, Issue 1, pp. 286, (2017) (PubMed).


Guo, Wang, Liu, Myatt, Sun: "Induction of PGF2? synthesis by cortisol through GR dependent induction of CBR1 in human amnion fibroblasts." in: **Endocrinology**, Vol. 155, Issue 8, pp. 3017-24, (2014) (PubMed).

Images




Flow Cytometry

Image 1. UBE3A Antibody (C-term) (ABIN388925 and ABIN2850432) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Western Blotting

Image 2. Western blot analysis of UBE3A Antibody (C-term) (ABIN388925 and ABIN2850432) in T47D cell line lysates (35 μ g/lane).UBE3A(arrow) was detected using the purified Pab.



Immunofluorescence

Image 3. Confocal immunofluorescent analysis of UBE3A Antibody (C-term) (ABIN388925 and ABIN2850432) with Hela cell followed by Alexa Fluor 488-conjugated goat antirabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 phalloidin (red).DI was used to stain the cell nuclear (blue).

Please check the product details page for more images. Overall 4 images are available for ABIN388925.