

Datasheet for ABIN389180
anti-DAAM1 antibody (C-Term)[Go to Product page](#)

2 Images

1 Publication

Overview

Quantity:	400 µL
Target:	DAAM1
Binding Specificity:	AA 1034-1063, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DAAM1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This DAAM1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1034-1063 amino acids from the C-terminal region of human DAAM1.
Clone:	RB10612
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	DAAM1
Alternative Name:	DAAM1 (DAAM1 Products)
Background:	Functions of the cell cortex, including motility, adhesion, and cytokinesis, are mediated by the

Target Details

reorganization of the actin cytoskeleton and recent evidence suggests a role for the Formin homology (FH) proteins in these processes. DAAM1 contains FH domains and belongs to a novel FH protein subfamily implicated in cell polarity. Wnt/Fz signaling activates the small GTPase Rho, a key regulator of cytoskeleton architecture, to control cell polarity and movement during development. Activation requires Dvl-Rho complex formation, an assembly mediated by DAAM1, which is thought to function as a scaffolding protein.

Molecular Weight: 123473

Gene ID: 23002

NCBI Accession: [NP_001257449](#), [NP_055807](#)

UniProt: [Q9Y4D1](#)

Pathways: [WNT Signaling](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:10~50

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C


Storage Comment: Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.

Expiry Date: 6 months

Publications


Product cited in: Wang, Liu, Zhao, Song, Wang, Wang, Sun: "Phosphorylation of CDK2 on threonine 160 influences silencing of sex chromosome during male meiosis." in: **Biology of reproduction**, Vol.

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human brain tissue reacted with DM1 Antibody (C-term) (ABIN389180 and ABIN2839343) , which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated.



Western Blotting

Image 2. Western blot analysis of DM1 (arrow) using rabbit polyclonal DM1 Antibody (Human C-term) (ABIN389180 and ABIN2839343). 293 cell lysates (2 µg/lane) either nontransfected (Lane 1) or transiently transfected with the DM1 gene (Lane 2) (Origene Technologies).