

Datasheet for ABIN389595

anti-c-MET antibody (pTyr1349)[2 Images](#)[2 Publications](#)[Go to Product page](#)

Overview

Quantity:	400 µL
Target:	c-MET (MET)
Binding Specificity:	pTyr1349
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This c-MET antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This MET Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y1349 of human MET.
Clone:	RB5215
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	c-MET (MET)
Alternative Name:	MET (MET Products)
Background:	The proto-oncogene MET product is the hepatocyte growth factor receptor and encodes

Target Details

tyrosine-kinase activity. The primary single chain precursor protein is post-translationally cleaved to produce the alpha and beta subunits, which are disulfide linked to form the mature receptor. Various mutations in the MET gene are associated with papillary renal carcinoma.

Molecular Weight: 155541

Gene ID: 4233

NCBI Accession: [NP_000236](#), [NP_001120972](#)

UniProt: [P08581](#)

Pathways: [RTK Signaling](#), [Carbohydrate Homeostasis](#), [Synaptic Membrane](#), [Signaling of Hepatocyte Growth Factor Receptor](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:50~100

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Storage Comment: Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.

Expiry Date: 6 months

Publications

Product cited in: Schwab, Sison, Meade, Broniowska, Corbett, Ebert: "Decreased Sirtuin Deacetylase Activity in LRRK2 G2019S iPSC-Derived Dopaminergic Neurons." in: **Stem cell reports**, Vol. 9, Issue 6, pp. 1839-1852, (2018) ([PubMed](#)).


Takumida, Takumida, Katagiri, Anniko: "Localization of sirtuins (SIRT1-7) in the aged mouse inner ear." in: **Acta oto-laryngologica**, pp. 1-12, (2015) ([PubMed](#)).

He, Hu, Shi, Weidert, Lu, Xu, Huang, Kelley, Xie: "Activation of the aryl hydrocarbon receptor sensitizes mice to nonalcoholic steatohepatitis by deactivating mitochondrial sirtuin deacetylase Sirt3." in: **Molecular and cellular biology**, Vol. 33, Issue 10, pp. 2047-55, (2013) ([PubMed](#)).

Kamarajan, Alhazzazi, Danciu, Dsilva, Verdin, Kapila: "Receptor-interacting protein (RIP) and Sirtuin-3 (SIRT3) are on opposite sides of anoikis and tumorigenesis." in: **Cancer**, Vol. 118, Issue 23, pp. 5800-10, (2012) ([PubMed](#)).


Parker, Vazquez-Manrique, Tourette, Farina, Offner, Mukhopadhyay, Orfila, Darbois, Menet, Tissenbaum, Neri: "Integration of β -catenin, sirtuin, and FOXO signaling protects from mutant huntingtin toxicity." in: **The Journal of neuroscience : the official journal of the Society for Neuroscience**, Vol. 32, Issue 36, pp. 12630-40, (2012) ([PubMed](#)).

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated. BC = breast carcinoma, HC = hepatocarcinoma.



Western Blotting

Image 2. The anti-MET- Pab (ABIN389595 and ABIN2839608) is used in Western blot to detect MET- in mouse kidney tissue lysate.