

Datasheet for ABIN389876 anti-p53 antibody (pThr55)



Overview

Target:

Alternative Name:

Background:



Quantity: 400 µL Target: p53 (TP53) Binding Specificity: pThr55 Human Reactivity: Rabbit Host: Clonality: Polyclonal Conjugate: This p53 antibody is un-conjugated Application: Dot Blot (DB) **Product Details** This p53 Antibody is generated from rabbits immunized with a KLH conjugated synthetic Immunogen: phosphopeptide corresponding to amino acid residues surrounding T55 of human p53. RB07990 Clone: Isotype: Ig Fraction Purification: This antibody is purified through a protein A column, followed by peptide affinity purification. **Target Details**

Tumor protein p53, a nuclear protein, plays an essential role in the regulation of cell cycle,

p53 (TP53)

p53 (TP53 Products)

| specifically in the transition from G0 to G1. It is found in very low levels in normal cells, however, |
|--|
| in a variety of transformed cell lines, it is expressed in high amounts, and believed to contribute |
| to transformation and malignancy. p53 is a DNA-binding protein containing DNA-binding, |
| oligomerization and transcription activation domains. It is postulated to bind as a tetramer to a |
| p53-binding site and activate expression of downstream genes that inhibit growth and/or |
| invasion, and thus function as a tumor suppressor. Mutants of p53 that frequently occur in a |
| number of different human cancers fail to bind the consensus DNA binding site, and hence |
| cause the loss of tumor suppressor activity. Alterations of the TP53 gene occur not only as |
| somatic mutations in human malignancies, but also as germline mutations in some cancer- |
| prone families with Li-Fraumeni syndrome. |
| |

| Molecular Weight: | 43653 |
|-------------------|--|
| Gene ID: | 7157 |
| NCBI Accession: | NP_000537, NP_001119584, NP_001119585, NP_001119586, NP_001119587, NP_001119588, NP_001119589, NP_001119590, NP_001263624, NP_001263625, NP_001263626, NP_001263627, NP_001263628, NP_00126 |
| UniProt: | P04637 |
| Pathways: | p53 Signaling, MAPK Signaling, PI3K-Akt Signaling, Apoptosis, AMPK Signaling, Chromatin Binding, ER-Nucleus Signaling, Positive Regulation of Endopeptidase Activity, Hepatitis C, Protein targeting to Nucleus, Autophagy, Warburg Effect |

Application Details


Application Notes:

DB: 1:500

| Restrictions: | For Research Use only |
|--------------------|--|
| Handling | |
| Format: | Liquid |
| Buffer: | Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |

Handling

| Storage Comment: | Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles. |
|------------------|--|
| Expiry Date: | 6 months |
| lmages | |



Dot Blot

Image 1. Dot blot analysis of anti-p53-pT55 Pab (R) on nitrocellulose membrane. 50 ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are $0.5 \, \mu g$ per ml.