

Datasheet for ABIN389913 anti-LC3B antibody (pThr12)

1 Image

1 Publication



Overview

Quantity:	400 µL
Target:	LC3B (MAP1LC3B)
Binding Specificity:	pThr12
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Dot Blot (DB)
Product Details	
Immunogen:	This LC3B Antibody is generated from rabbits immunized with a KLH conjugated synthetic
	phosphopeptide corresponding to amino acid residues surrounding T12 of human LC3B.
Clone:	RB12679
Isotype:	Ig Fraction
Predicted Reactivity:	В
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target [.]	LC3B (MAP1LC3B)

Target:	LC3B (MAP1LC3B)
Alternative Name:	LC3B (MAP1LC3B Products)
Background:	MAP1A and MAP1B are microtubule-associated proteins which mediate the physical

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN389913 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Molecular Weight:	14688
	degraded within the lysosome (or vacuole).
	lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then
	constituent targeted for degradation in a membrane bound structure, which then fuse with the
	formation of double-membrane bound autophagosomes which enclose the cytoplasmic
	cytoplasmic enzymes and organelles during nutrient starvation. Macroautophagy involves the
	cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active
	bound form, LC3-II. Macroautophagy is the major inducible pathway for the general turnover of
	APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form the membrane-
	molecule is cleaved by APG4B/ATG4B to form the cytosolic form, LC3-I. This is activated by
	the light chain subunits and can associate with either MAP1A or MAP1B. The precursor
	each consist of a heavy chain subunit and multiple light chain subunits. MAP1LC3b is one of
	involved in formation of autophagosomal vacuoles (autophagosomes). MAP1A and MAP1B
	interactions between microtubules and components of the cytoskeleton. These proteins are

Gene ID:	81631
NCBI Accession:	NP_073729
UniProt:	Q9GZQ8
Pathways:	Autophagy

Application Details

Application Notes:	DB: 1:500
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN389913 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Handling

	aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months
Publications	
Product cited in:	Mitroulis, Kourtzelis, Kambas, Rafail, Chrysanthopoulou, Speletas, Ritis: "Regulation of the
	autophagic machinery in human neutrophils." in: European journal of immunology, Vol. 40,
	lssue 5, pp. 1461-72, (2010) (PubMed).

Images



Dot Blot

Dot Blot

Image 1. Dot blot analysis of Phospho-LC3 (G8b)- T12 Antibody 3530a on nitrocellulose membrane. 50 ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are $0.5 \,\mu g$ per ml.