antibodies - online.com







anti-SLAMF1 antibody (pTyr281)







()	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	$I \vee I$	ew

Quantity:	400 μL
Target:	SLAMF1
Binding Specificity:	pTyr281
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLAMF1 antibody is un-conjugated
Application:	Dot Blot (DB)

Product Details

Immunogen:	This SLAMF1 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y281 of human SLAMF1.
Clone:	RB15251
Isotype:	lg Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	SLAMF1
Alternative Name:	SLAMF1 (SLAMF1 Products)
Background: SLAMF1 is a high-affinity self-ligand important in bidirectional T-cell to B-cell stimulation. Sl	

Target Details

	induced signal-transduction events in T-lymphocytes are different from those in B-cells. Two	
	modes of SLAM signaling are likely to exist: one in which the inhibitor SH2D1A acts as a	
	negative regulator and another in which protein-tyrosine phosphatase 2C (PTPN11)-dependent	
	signal transduction operates.	
Molecular Weight:	37231	
Gene ID:	6504	
NCBI Accession:	NP_003028	

Application Details


Q13291

Application Notes:	DB: 1:500
Restrictions:	For Research Use only

Handling

UniProt:

Format:	Liquid	
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.	
Expiry Date:	6 months	



Dot Blot

Image 1. Dot blot analysis of anti-Phospho-SLF1-p Antibody (ABIN389924 and ABIN2839746) on nitrocellulose membrane. 50 ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5 μg per ml.