

Datasheet for ABIN389970
anti-TNK2 antibody (pTyr518)



[Go to Product page](#)

1 Image

Overview

Quantity:	400 µL
Target:	TNK2
Binding Specificity:	pTyr518
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TNK2 antibody is un-conjugated
Application:	Dot Blot (DB)

Product Details

Immunogen:	This ACK1 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y518 of human ACK1.
Clone:	RB15592
Isotype:	Ig Fraction
Predicted Reactivity:	B
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	TNK2
Alternative Name:	ACK1 (TNK2 Products)

Target Details


Background:	ACK1 is a tyrosine kinase that binds Cdc42Hs in its GTP-bound form and inhibits both the intrinsic and GTPase-activating protein (GAP)-stimulated GTPase activity of Cdc42Hs. This binding is mediated by a unique sequence of 47 amino acids C-terminal to an SH3 domain. The protein may be involved in a regulatory mechanism that sustains the GTP-bound active form of Cdc42Hs and which is directly linked to a tyrosine phosphorylation signal transduction pathway.
Molecular Weight:	114569
Gene ID:	10188
NCBI Accession:	NP_001010938 , NP_005772
UniProt:	Q07912

Application Details

Application Notes:	DB: 1:500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months



Dot Blot

Image 1. Dot blot analysis of anti-Phospho-ACK1-p Antibody (ABIN389970 and ABIN2839765) on nitrocellulose membrane. 50 ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5 μ g per ml.