antibodies -online.com





anti-ACE2 antibody (C-Term)





Publication



Go to Product page

_					
	W	0	rv	10	W

Alternative Name:

Background:

Quantity:	400 μL
Target:	ACE2
Binding Specificity:	AA 773-805, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This ACE2 (SARS Receptor) antibody is generated from rabbits immunized with a KLH
	conjugated synthetic peptide between 773-805 amino acids from the C-terminal region of human ACE2 (SARS Receptor).
Clone:	RB4819
Isotype:	Ig Fraction
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by
	dialysis against PBS.
Target Details	
Target:	ACE2

ACE2 cDNA encodes a deduced 805-amino acid protein containing a potential 17-amino acid N-

ACE2 (SARS Receptor) (ACE2 Products)

terminal signal peptide and a putative 22-amino acid C-terminal membrane anchor. It also possesses a zinc metalloprotease consensus sequence and a conserved glutamine residue that may function as a third zinc ligand. ACE2 is expressed predominantly in vascular endothelial cells of the heart and kidney. ACE converts angiotensin I to angiotensin II, ACE2 converts angiotensin I to angiotensin I to angiotensin I -9, which has 9 amino acids. Angiotensin II is a potent blood vessel constrictor, while angiotensin 1-9 does not impact blood vessels but is cleaved by ACE to a shorter peptide, angiotensin 1-7, which is a blood vessel dilator. Spike (S) proteins of coronaviruses, including the SARS coronavirus, bind with cellular receptors to mediate infection of target cells. ACE2 binds the S1 domain of the SARS coronavirus S protein. SARS coronavirus replicates efficiently on ACE2-transfected but not mock-transfected 293T cells. Anti-ACE2 but not anti-ACE1 antibody blocks viral replication on Vero E6 cells. It has been proposed that ACE2 is a functional receptor for SARS coronavirus.

Molecular Weight:	92463
Gene ID:	59272
NCBI Accession:	NP_068576
UniProt:	Q9BYF1
Pathways:	ACE Inhibitor Pathway, Peptide Hormone Metabolism, Regulation of Systemic Arterial Blood
	Pressure by Hormones, Feeding Behaviour

Application Details

Application Notes:	WB: 1:1000. WB: 1:1000. IHC-P: 1:50~100. IHC-P: 1:50~100
Restrictions:	For Research Use only

Handling

Format:	Liquid	
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in sma	

aliquots to prevent freeze-thaw cycles.

Expiry Date:


6 months


Publications

Product cited in:

Heurich, Hofmann-Winkler, Gierer, Liepold, Jahn, Pöhlmann: "TMPRSS2 and ADAM17 cleave ACE2 differentially and only proteolysis by TMPRSS2 augments entry driven by the severe acute respiratory syndrome coronavirus spike protein." in: **Journal of virology**, Vol. 88, Issue 2, pp. 1293-307, (2013) (PubMed).

Images






Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human testis tissue reacted with ACE2 (SARS Receptor) Antibody (C-term) f , which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated.

Western Blotting

Image 2. The anti-ACE2 C-term Pab f is used in Western blot to detect ACE2 in 293 cell lysate.



Western Blotting

Image 3. Western blot analysis of anti-ACE2 C-term Pab f in K562 cell line lysates (35 μ g/lane). ACE2(arrow) was detected using the purified Pab.

Please check the product details page for more images. Overall 4 images are available for ABIN390049.