

Datasheet for ABIN390215

anti-CDC25A antibody (AA 57-84)[Go to Product page](#)**1** Image

Overview

Quantity:	400 µL
Target:	CDC25A
Binding Specificity:	AA 57-84
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CDC25A antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	This Cdc25A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 57-84 amino acids from human Cdc25A.
Clone:	RB08125
Isotype:	Ig Fraction
Predicted Reactivity:	B, M, Rat
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	CDC25A
Alternative Name:	Cdc25A (CDC25A Products)

Target Details


Background:	CDC25A is a member of the CDC25 family of phosphatases. CDC25A is required for progression from G1 to the S phase of the cell cycle. It activates the cyclin-dependent kinase CDC2 by removing two phosphate groups. CDC25A is specifically degraded in response to DNA damage, which prevents cells with chromosomal abnormalities from progressing through cell division. CDC25A is an oncogene, although its exact role in oncogenesis has not been demonstrated.
Molecular Weight:	59087
Gene ID:	993
NCBI Accession:	NP_001780 , NP_963861
UniProt:	P30304
Pathways:	Cell Division Cycle , Mitotic G1-G1/S Phases , M Phase

Application Details

Application Notes:	WB: 1:1000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months



Western Blotting

Image 1. Cdc25A Antibody (S78) (ABIN390215 and ABIN2840697) western blot analysis in MCF-7 cell line lysates (35 µg/lane). This demonstrates the Cdc25A antibody detected the Cdc25A protein (arrow).