

Datasheet for ABIN390253
anti-NR4A1 antibody (AA 329-358)[Go to Product page](#)**1** Image**1** Publication

Overview

Quantity:	400 µL
Target:	NR4A1
Binding Specificity:	AA 329-358
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NR4A1 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	This NUR77(NR4A1) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 329-358 amino acids from human NUR77(NR4A1).
Clone:	RB11093
Isotype:	Ig Fraction
Predicted Reactivity:	M, Rat
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	NR4A1
Alternative Name:	NUR77 (NR4A1) (NR4A1 Products)

Target Details

Background:	NR4A1 is a member of the steroid-thyroid hormone-retinoid receptor superfamily. Expression is induced by phytohemagglutinin in human lymphocytes and by serum stimulation of arrested fibroblasts. This protein acts as a nuclear transcription factor. Translocation of the protein from the nucleus to mitochondria induces apoptosis.
Molecular Weight:	64463
Gene ID:	3164
NCBI Accession:	NP_001189162 , NP_002126 , NP_775180
UniProt:	P22736
Pathways:	Fc-epsilon Receptor Signaling Pathway , Nuclear Receptor Transcription Pathway , EGFR Signaling Pathway , Neurotrophin Signaling Pathway , Steroid Hormone Mediated Signaling Pathway

Application Details

Application Notes:	WB: 1:500
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Publications

Product cited in:	Xu, Han, Epstein, Liu: "Regulation of PDK mRNA by high fatty acid and glucose in pancreatic islets." in: Biochemical and biophysical research communications , Vol. 344, Issue 3, pp. 827-
-------------------	---

Images



Western Blotting

Image 1. All lanes : Anti-NUR77 (NR4A1) Antibody at 1:500 dilution Lane 1: Jurkat whole cell lysate Lane 2: HeLa whole cell lysate Lane 3: Human liver tissue lysate Lane 4: MCF-7 whole cell lysate Lane 5: Mouse brain tissue lysate Lane 6: NIH/3T3 whole cell lysate Lane 7: Rat brain tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 64 kDa Blocking/Dilution buffer: 5 % NFDm/TBST.