# antibodies - online.com







## anti-Frataxin antibody (C-Term)

**Images** 




Go to	D		
	Pron	ויאווו	Dage

Overview	
Quantity:	400 μL
Target:	Frataxin (FXN)
Binding Specificity:	AA 143-172, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Frataxin antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This FXN antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 143-172 amino acids from the C-terminal region of human FXN.
Clone:	RB12571
Isotype:	lg Fraction
Predicted Reactivity:	М
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Target Details	
Target:	Frataxin (FXN)


### **Target Details**

raiget Details	
Alternative Name:	FXN (FXN Products)
Background:	FXN is a mitochondrial protein which belongs to the FRATAXIN family. The protein functions in regulating mitochondrial iron transport and respiration. The expansion of intronic trinucleotide repeat GAA results in Friedreich ataxia.
Molecular Weight:	23135
Gene ID:	2395
NCBI Accession:	NP_000135, NP_001155178, NP_852090
UniProt:	Q16595
Pathways:	Transition Metal Ion Homeostasis, Regulation of Cell Size
Application Details	
Application Notes:	WB: 1:1000. WB: 1:1000. IHC-P: 1:10~50
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months




#### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with FXN antibody (C-term) (ABIN390377 and ABIN2840779), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated.



#### **Western Blotting**

**Image 2.** Western blot analysis of anti-FXN Antibody (Cterm) (ABIN390377 and ABIN2840779) in 293 cell line lysates (35  $\mu$ g/lane). FXN (arrow) was detected using the purified Pab.



#### **Western Blotting**

**Image 3.** Western blot analysis of FXN (arrow) using rabbit polyclonal FXN Antibody (C-term) (R). 293 cell lysates (2  $\mu$  g/lane) either nontransfected (Lane 1) or transiently transfected with the FXN gene (Lane 2) (Origene Technologies).